

Random Variables Intro II

Expectation: just like a weighted average; we weight the values that X can take on by the probabilities of getting those values. Expectation is defined as

$$\mathbb{E}[X] = \sum_k k \cdot \mathbb{P}[X = k].$$

If X is a non-negative integer-valued random variable, we have the *tail sum formula*:

$$\mathbb{E}[X] = \sum_{k \geq 1} \mathbb{P}[X \geq k].$$

The *Law of the Unconscious Statistician (LOTUS)*:

$$\mathbb{E}[f(X)] = \sum_k f(k) \cdot \mathbb{P}[X = k].$$

Linearity of Expectation: for two random variables X, Y (which could be dependent),

$$\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y].$$

Joint Distributions

With two RVs, their *joint distribution* is $\mathbb{P}[X = x, Y = y]$. The *marginal distributions* are

$$\mathbb{P}[X = x] = \sum_y \mathbb{P}[X = x, Y = y] \quad \mathbb{P}[Y = y] = \sum_x \mathbb{P}[X = x, Y = y]$$

The conditional probability with two random variables is defined as

$$\mathbb{P}[X = x \mid Y = y] = \frac{\mathbb{P}[X = x, Y = y]}{\mathbb{P}[Y = y]}.$$

Two random variables are *independent* if and only if

$$\mathbb{P}[X = x, Y = y] = \mathbb{P}[X = x] \mathbb{P}[Y = y] \quad \mathbb{P}[X = x \mid Y = y] = \mathbb{P}[X = x].$$

1 Pullout Balls

Note 15

Suppose you have a bag containing four balls numbered 1, 2, 3, 4.

(a) You perform the following experiment: pull out a single ball and record its number. What is the expected value of the number that you record?

(b) You repeat the experiment from part (a), except this time you pull out two balls together and record the product of their numbers. What is the expected value of the total that you record?

2 Linearity

Note 15

Solve each of the following problems using linearity of expectation. Explain your methods clearly.

- (a) In an arcade, you play game A 10 times and game B 20 times. Each time you play game A , you win with probability $1/3$ (independently of the other times), and if you win you get 3 tickets (redeemable for prizes), and if you lose you get 0 tickets. Game B is similar, but you win with probability $1/5$, and if you win you get 4 tickets. What is the expected total number of tickets you receive?
- (b) A monkey types at a 26-letter keyboard with one key corresponding to each of the lower-case English letters. Each keystroke is chosen independently and uniformly at random from the 26 possibilities. If the monkey types 1 million letters, what is the expected number of times the sequence “book” appears? (*Hint*: Consider where the sequence “book” can appear in the string.)

3 Family Planning

Note 15

Mr. and Mrs. Johnson decide to continue having children until they either have their first girl or until they have three children. Assume that each child is equally likely to be a boy or a girl, independent of all other children, and that there are no multiple births. Let G denote the numbers of girls that the Johnsons have. Let C be the total number of children they have.

(a) Determine the sample space, along with the probability of each sample point.

(b) Compute the joint distribution of G and C . Fill in the table below.

	$C = 1$	$C = 2$	$C = 3$
$G = 0$			
$G = 1$			

(c) Use the joint distribution to compute the marginal distributions of G and C and confirm that the values are as you'd expect. Fill in the tables below.

$\mathbb{P}[G = 0]$		$\mathbb{P}[C = 1]$	$\mathbb{P}[C = 2]$	$\mathbb{P}[C = 3]$
$\mathbb{P}[G = 1]$				

(d) Are G and C independent?

(e) What is the expected number of girls the Johnsons will have? What is the expected number of children that the Johnsons will have?