#### Counting – Part 1

#### CS70: Discrete Mathematics and Probability Theory

#### UC Berkeley – Summer 2025

Lecture 12

Ref: Note 10

# Probability

Wait! Probably doesn't come until later! A little preview: A bag contains:

What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide.

Today: Counting!

Later: Probability.

# A Look Into The Future!

What's to come? Probability.

A bag contains:



What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide. Chances?

- (A) Red probability is 3/8?
- (B) Blue probability is 3/9?
- (C) Blue probability is 3/8?
- (D) Yellow probability is 2/8?

You can all count!

Combinatorics is fancy counting.



- 2 Tree of Choices
- Rules of Counting
- Sample with/without replacement where order does/doesn't matter

- How many outcomes possible for k coin tosses?
- How many poker hands?
- How many handshakes for *n* people?
- How many diagonals in a *n* sided convex polygon?
- How many 10 digit numbers?
- How many 10 digit numbers without repetition?
- How many ways can I divide up 5 dollars among 3 people?

How many 3-bit binary strings? How many different sequences of three bits from {0,1}? How would you make one sequence? How many different ways to do that making?



8 leaves which is  $2 \times 2 \times 2$ . One leaf for each string.

 $\Rightarrow$  8 3-bit strings!

#### First Rule of Counting: Product Rule

Objects made by choosing from  $n_1$ , then from  $n_2, ...,$  then from  $n_k$  ... the number of objects is  $n_1 \times n_2 \cdots \times n_k$ .



In picture,  $2 \times 2 \times 3 = 12$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \times \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

```
10 ways for first choice, 10 ways for second choice, ... 10 \times 10 \times \cdots \times 10 = 10^{10}
```

How many *n* digit base *m* numbers? *m* ways for first, *m* ways for second, ...  $m \times m \times \cdots \times m = m^n$ 

But: Is 09, a two digit number? If not, then  $(m-1)m^{n-1}$ 

# **Counting Functions and Polynomials**

How many functions *f* mapping *S* to *T*? (i.e.,  $f : S \rightarrow T$ )

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ... ...  $|T|^{|S|}$ 

How many polynomials of degree d modulo p?

*p* ways to choose for first coefficient, *p* ways for second, ... ... *p*<sup>*d*+1</sup>

Counting by points at x = 0, 1, 2, ..., d rather than coefficients (d + 1 points!):

p values for first y value, p values for second, ...  $\dots p^{d+1}$ 

Questions?

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

 $\dots 10 \times 9 \times 8 \times \dots \times 1 = 10!$ 

Sample of size *k* from *n* numbers without replacement? *n* ways for first choice, n-1 ways for second, n-2 choices for third, ... ...  $n \times (n-1) \times (n-2) \times \cdots \times (n-k+1) = \frac{n!}{(n-k)!}$ 

How many orderings of *n* objects? (**Permutations of** *n* **objects**) *n* ways for first, n-1 ways for second, n-2 ways for third, ...

 $\dots n \times (n-1) \times (n-2) \times \dots \times 1 = n!$ 

How many one-to-one functions from S to S (bijections on S)?

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

 $\dots |S| \times (|S|-1) \times \dots 1 = |S|!$ 

A one-to-one function with the same domain and range is a permutation!

Bijections are permutations – recall:

 $f(x) = 2x \pmod{5}$   $[0, 1, 2, 3, 4] \rightarrow [0, 2, 4, 1, 3]$ 

Number of bijections from  $\{0, 1, 2, 3, 4\}$  to  $\{0, 1, 2, 3, 4\}$ ?

Number of bijections like  $f(x) = ax \pmod{5}$ ?

Linear bijections:  $f(x) = ax + b \pmod{5}$ ?

 $\implies$  linear functions (mod p) are a tiny fraction of bijections on  $\{0, \dots, p-1\}$ 

#### Counting Sets – When Order Doesn't Matter

How many poker hands? (52 cards in a deck, all different, 5 in a poker hand)

 $52 \times 51 \times 50 \times 49 \times 48 = \frac{52!}{47!}$ ???

Are A, K, Q, 10, J of spades and 10, J, Q, K, A of spades the same?

Second Rule of Counting: If order doesn't matter, count ordered objects and then divide by the number of equivalent orderings of an ordered object.

"the" number of equivalent... assumes same for all ordered objects!

Number of orderings for a poker hand: 5!

By second rule of counting, number of poker hands is

$$\frac{52!}{47!} \cdot \frac{1}{5!} = \frac{52!}{47! \cdot 5!}$$

Generic: ways to choose 5 out of 52 possibilities.

### Ordered to Unordered

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by the number of equivalent orderings.



How many red nodes (ordered objects)? 9

How many red nodes mapped to one blue node? 3

How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ 

How many poker deals? 52.51.50.49.48

How many poker deals per hand? Map each deal to ordered deal: 5! How many poker hands?  $\frac{52 \cdot 51 \cdot 50 \cdot 49 \cdot 48}{5!}$ Questions?

#### When Order Doesn't Matter

Choose 2 out of n?

$$\frac{n\times(n-1)}{2} = \frac{n!}{(n-2)!\times 2}$$

Choose 3 out of n?

$$\frac{n \times (n-1) \times (n-2)}{3!} = \frac{n!}{(n-3)! \times 3!}$$

Choose k out of n?

 $\frac{n!}{(n-k)! \times k!}$ 

Notation:  $\binom{n}{k}$  and pronounced "*n* choose *k*" Familiar? Questions?

## Example: Visualize the Proof

*k* choices. *First Rule:*  $n_1 \times \cdots \times n_k$ *Second Rule:* when order doesn't matter, divide...



3 card poker deals:  $52 \times 51 \times 50 \ (=\frac{52!}{49!})$ . First Rule

Poker hands:  $\Delta$ ? Hand: Q, K, A. Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K  $\Delta = 3 \times 2 \times 1$  First Rule again Total:  $\frac{52!}{49!3!}$  Second Rule! Choose k out of n: Ordered set:  $\frac{n!}{(n-k)!}$  (First Rule) Orderings of one hand? k! (First Rule)  $\implies$  Total:  $\frac{n!}{(n-k)!k!}$  (Second rule)

### Example: Anagrams

*k* choices. *First Rule:*  $n_1 \times \cdots \times n_k$ *Second Rule:* when order doesn't matter, divide...



For each of these: correct or not?

- (A) Number of poker hands  $= \binom{52}{5}$
- (B) Orderings of "CAT" = 3!
- (C) Orderings of ANAGRAM =  $\frac{7!}{3!}$
- (D) Orderings of ANAGRAM =  $\frac{7!}{4!}$
- (E) Orders of MISSISSIPPI =  $\frac{11!}{4!4!2!}$
- (F) Orders of MISSISSIPPI =  $\frac{11!}{10!}$

### Explanations

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3 × 2 × 1 = 3! orderings

How many orderings of the letters in ANAGRAM?

```
Ordered, except for A
```

```
Total orderings of 7 letters = 7!
```

```
Total equivalent orderings of three A's = 3!
```

```
Total orderings? 7!
```

How many orderings of MISSISSIPPI?

11 letters total: 4 S's, 4 I's, 2 P's

11! ordered objects.

 $4! \times 4! \times 2!$  ordered objects per "unordered object"

Total orderings? 11!

*Important:* Not just 10 letters rearranged – 3 groups of letters Choose order of S's (4!), choose order of I's (4!), choose order of P's (2!). Multiply (*First Rule*)

### How To Sample k Items Out Of n

Without replacement:

Order matters: 
$$n \times (n-1) \times (n-2) \dots \times (n-k+1) = \frac{n!}{(n-k)!}$$

Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}$  or "*n* choose *k*" or  $\binom{n}{k}$ 

With Replacement:

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

Problem: depends on how many of each item we chose! Different number of ordered items map to each unordered item

| Jnordered item: 1,2,3 | 3! ordered items map to              |
|-----------------------|--------------------------------------|
| Jnordered item: 1,2,2 | $\frac{3!}{2!}$ ordered items map to |

How do we deal with this mess??

it it

# Splitting Up Money: The Problem

Problem: How many ways can Bob and Alice split 5 dollars?

For each of 5 dollars pick Bob or Alice (2<sup>5</sup>), divide out order???

5 dollars for Bob and 0 for Alice: one ordered set: (*B*, *B*, *B*, *B*, *B*)

4 for Bob and 1 for Alice:

5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...

"Sorted" way to specify, first Alice's dollars, then Bob's. (B, B, B, B, B): 1: (B,B,B,B,B) (A, B, B, B, B): 5: (A,B,B,B,B), (B,A,B,B,B), (B,B,A,B,B),... (A, A, B, B, B):  $\binom{5}{2}$ : (A,A,B,B,B), (A,B,A,B,B), (A,B,B,A,B), ... and so on.



Second rule of counting is no good here!

# Splitting Up Money: Stars and Bars

Problem: How many ways can Alice, Bob, and Eve split 5 dollars?

One way: Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E)

Idea: Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars:  $\star \star \star \star$ Separate with bars: Two "middle separators" (bars) for three sets.  $(A, A, A, B, E) \longrightarrow$  Stars and Bars:  $\star \star \star | \star | \star$ .

Alice: 2, Bob: 1, Eve: 2.

 $\rightarrow$  Stars and Bars: \*\*|\*|\*\*.

Alice: 0, Bob: 1, Eve: 4.

 $\longrightarrow$  Stars and Bars:  $|\star| \star \star \star \star$ .

Each split "is" a sequence of stars and bars

... and each sequence of stars and bars "is" a split

... a bijection!

# Counting Rule: If there is a bijection between two sets they have the same size!

Question: How many different 5 star and 2 bar diagrams?

```
Regardless of how split: 7 symbols – 5 are stars, 2 are bars
To help visualize: Number the symbol positions 1 to 7
```

```
Alice: 0; Bob 1; Eve: 4

\begin{vmatrix} * & | & * & * & * \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{vmatrix}
Bars in positions 1 and 3

Alice: 1; Bob 4; Eve: 0

\begin{vmatrix} * & | & * & * & * \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{vmatrix}
Bars in positions 2 and 7
```

Picking 2 positions for bars ... out of 7 positions ...

 $\binom{7}{2}$  ways to pick  $\longrightarrow \binom{7}{2}$  ways to split 5 dollars among 3 people.

#### Stars and Bars

Back to general problem:

Sample *k* from *n* with replacement, where order doesn't matter  $\Rightarrow$  Only the count of each of the *n* items matters

k selections (stars); separated into n groups with n-1 bars

\*\* \* ··· \*\*

k stars, n-1 bars  $\longrightarrow n+k-1$  length string

Select n-1 of the n+k-1 positions for the bars

Ways to do so:

 $\binom{n+k-1}{n-1}$ 

Could also select positions for the k stars – gives the same count:

 $\binom{n+k-1}{k}$ 

For each of these: correct or not?

- (A) Ways to split *k* dollars among *n*:  $\binom{k+n-1}{n-1}$
- (B) Ways to split *n* dollars among k:  $\binom{n+k-1}{k-1}$
- (C) Ways to split 5 dollars among 3:  $\binom{5+3-1}{3-1}$
- (D) Ways to split 5 dollars among 3:  $\binom{7}{5}$

**First rule:**  $n_1 \times n_2 \times \cdots \times n_k$ 

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

#### Second rule: when order doesn't matter divide .. when possible

Sample without replacement, no order:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

#### One-to-one rule: equal in number if one-to-one correspondence

Sample with replacement and order doesn't matter:  $\binom{k+n-1}{n-1}$ Distribute *k* samples (stars) over *n* poss. (*n*-1 bars group poss..) Distribute *k* dollars to *n* people.