

## 1 Proof Practice

- (a) Prove that  $\forall n \in \mathbb{N}$ , if  $n$  is odd, then  $n^2 + 1$  is even. (Recall that  $n$  is odd if  $n = 2k + 1$  for some natural number  $k$ .)
- (b) Prove that  $\forall x, y \in \mathbb{R}$ ,  $\min(x, y) = (x + y - |x - y|)/2$ . (Recall, that the definition of absolute value for a real number  $z$ , is

$$|z| = \begin{cases} z, & z \geq 0 \\ -z, & z < 0 \end{cases}$$

- (c) Suppose  $A \subseteq B$ . Prove  $\mathcal{P}(A) \subseteq \mathcal{P}(B)$ . (Recall that  $A' \in \mathcal{P}(A)$  if and only if  $A' \subseteq A$ .)

### Solution:

- (a) We will use a direct proof. Assume  $n$  is odd. By the definition of odd numbers,  $n = 2k + 1$  for some natural number  $k$ . Substituting into the expression  $n^2 + 1$ , we get  $(2k + 1)^2 + 1$ . Simplifying the expression yields  $4k^2 + 4k + 2$ . This can be rewritten as  $2 \times (2k^2 + 2k + 1)$ . Since  $2k^2 + 2k + 1$  is a natural number, by the definition of even numbers,  $n^2 + 1$  is even.
- (b) We will use a proof by cases. Again, the definition of the absolute value function for real number  $z$  is

$$|z| = \begin{cases} z, & z \geq 0 \\ -z, & z < 0 \end{cases}$$

**Case 1:**  $x < y$ . This means  $|x - y| = y - x$ . Substituting this into the formula on the right hand side, we get

$$\frac{x + y - y + x}{2} = x = \min(x, y).$$

**Case 2:**  $x \geq y$ . This means  $|x - y| = x - y$ . Substituting this into the formula on the right hand side, we get

$$\frac{x + y - x + y}{2} = y = \min(x, y).$$

- (c) Suppose  $A' \in \mathcal{P}(A)$ , that is,  $A' \subseteq A$  (by the definition of the power set).

Let  $x \in A'$ . Then, since  $A' \subseteq A$ ,  $x \in A$ . Since  $A \subseteq B$ ,  $x \in B$ . We have shown  $(\forall x \in A') x \in B$ , so  $A' \subseteq B$ .

Since the previous argument works for any  $A' \subseteq A$ , we have proven  $(\forall A' \in \mathcal{P}(A)) A' \subseteq B$ . So,  $(\forall A' \in \mathcal{P}(A))(A' \in \mathcal{P}(B))$ . Thus, we conclude  $\mathcal{P}(A) \subseteq \mathcal{P}(B)$  as desired.

## 2 Preserving Set Operations

For a function  $f$ , define the image of a set  $X$  to be the set  $f(X) = \{y \mid y = f(x) \text{ for some } x \in X\}$ . Define the inverse image or preimage of a set  $Y$  to be the set  $f^{-1}(Y) = \{x \mid f(x) \in Y\}$ . Prove the following statements, in which  $A$  and  $B$  are sets. By doing so, you will show that inverse images preserve set operations, but images typically do not.

*Recall: For sets  $X$  and  $Y$ ,  $X = Y$  if and only if  $X \subseteq Y$  and  $Y \subseteq X$ . To prove that  $X \subseteq Y$ , it is sufficient to show that  $(\forall x) ((x \in X) \implies (x \in Y))$ .*

(a)  $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$ .

(b)  $f(A \cup B) = f(A) \cup f(B)$ .

### Solution:

In order to prove equality  $A = B$ , we need to prove that  $A$  is a subset of  $B$ ,  $A \subseteq B$  and that  $B$  is a subset of  $A$ ,  $B \subseteq A$ . To prove that LHS is a subset of RHS we need to prove that if an element is a member of LHS then it is also an element of the RHS.

(a) Suppose  $x \in f^{-1}(A \cup B)$  which means that  $f(x) \in A \cup B$ . Then either  $f(x) \in A$ , in which case  $x \in f^{-1}(A)$ , or  $f(x) \in B$ , in which case  $x \in f^{-1}(B)$ , so in either case we have  $x \in f^{-1}(A) \cup f^{-1}(B)$ . This proves that  $f^{-1}(A \cup B) \subseteq f^{-1}(A) \cup f^{-1}(B)$ .

Now, suppose that  $x \in f^{-1}(A) \cup f^{-1}(B)$ . Suppose, without loss of generality, that  $x \in f^{-1}(A)$ . Then  $f(x) \in A$ , so  $f(x) \in A \cup B$ , so  $x \in f^{-1}(A \cup B)$ . The argument for  $x \in f^{-1}(B)$  is the same. Hence,  $f^{-1}(A) \cup f^{-1}(B) \subseteq f^{-1}(A \cup B)$ .

(b) Suppose that  $x \in A \cup B$ . Then either  $x \in A$ , in which case  $f(x) \in f(A)$ , or  $x \in B$ , in which case  $f(x) \in f(B)$ . In either case,  $f(x) \in f(A) \cup f(B)$ , so  $f(A \cup B) \subseteq f(A) \cup f(B)$ .

Now, suppose that  $y \in f(A) \cup f(B)$ . Then either  $y \in f(A)$  or  $y \in f(B)$ . In the first case, there is an element  $x \in A$  with  $f(x) = y$ ; in the second case, there is an element  $x \in B$  with  $f(x) = y$ . In either case, there is an element  $x \in A \cup B$  with  $f(x) = y$ , which means that  $y \in f(A \cup B)$ . So  $f(A) \cup f(B) \subseteq f(A \cup B)$ .

The purpose of this problem is to gain familiarity to naming things precisely. In particular, we named an element in the LHS (or the pre-image of the LHS) and then argued about whether that element or its image was in the right hand side. By explicitly naming an element generically where it could be *any element in the set*, we could argue about its membership in a set and or its image or preimage. With these different concepts floating around it is helpful to be clear in the argument.

## 3 Pebbles

Suppose you have a rectangular array of pebbles, where each pebble is either red or blue. Suppose that for every way of choosing one pebble from each column, there exists a red pebble among the chosen ones. Prove that there must exist an all-red column.

**Solution:** We give a proof by contraposition. Suppose there does not exist an all-red column. This means that, in each column, we can find a blue pebble. Therefore, if we take one blue pebble from each column, we have a way of choosing one pebble from each column without any red pebbles. This is the negation of the original hypothesis, so we are done.