

1 Continuous Joint Densities

The joint probability density function of two random variables X and Y is given by $f(x,y) = Cxy$ for $0 \leq x \leq 1, 0 \leq y \leq 2$, and 0 otherwise (for a constant C).

(a) Find the constant C that ensures that $f(x,y)$ is indeed a probability density function.

(b) Find $f_X(x)$, the marginal distribution of X .

(c) Find the conditional distribution of Y given $X = x$.

(d) Are X and Y independent?

2 Uniform Distribution

You have two fidget spinners, each having a circumference of 10. You mark one point on each spinner as a needle and place each of them at the center of a circle with values in the range $[0, 10)$ marked on the circumference. If you spin both (independently) and let X be the position of the first spinner's mark and Y be the position of the second spinner's mark, what is the probability that $X \geq 5$, given that $Y \geq X$?

3 Exponential Practice

- (a) Let $X_1, X_2 \sim \text{Exponential}(\lambda)$ be independent, $\lambda > 0$. Calculate the density of $Y := X_1 + X_2$.
[*Hint*: One way to approach this problem would be to compute the CDF of Y and then differentiate the CDF.]

- (b) Let $t > 0$. What is the density of X_1 , conditioned on $X_1 + X_2 = t$? [*Hint*: Once again, it may be helpful to consider the CDF $\mathbb{P}(X_1 \leq x \mid X_1 + X_2 = t)$. To tackle the conditioning part, try conditioning instead on the event $\{X_1 + X_2 \in [t, t + \varepsilon]\}$, where $\varepsilon > 0$ is small.]