Message "is" points on P(x). (degree n-1, n+2k points.)

Message "is" points on P(x). (degree n-1, n+2k points.) Channel: Send P(i), receive R(i).

Message "is" points on P(x). (degree n-1, n+2k points.) Channel: Send P(i), receive R(i). Errors are wrong values at  $\leq k$  points.

Message "is" points on P(x). (degree n-1, n+2k points.) Channel: Send P(i), receive R(i). Errors are wrong values at  $\leq k$  points.Error:  $P(i) \neq R(i)$ .

Message "is" points on P(x). (degree n-1, n+2k points.) Channel: Send P(i), receive R(i).

Errors are wrong values at  $\leq k$  points.Error:  $P(i) \neq R(i)$ .

Error locator polynomial:  $E(x) = (x - e_1) \cdot (x - e_k) = x^k + b_{k-1}x^{k-1} + \dots + b_0.$ 

Message "is" points on P(x). (degree n-1, n+2k points.) Channel: Send P(i), receive R(i).

Errors are wrong values at  $\leq k$  points.Error:  $P(i) \neq R(i)$ .

Error locator polynomial:  $E(x) = (x - e_1) \cdot (x - e_k) = x^k + b_{k-1}x^{k-1} + \dots + b_0.$ Find:  $Q(x) = E(x)P(x) = a_{n+k-1}x^{n+k-1} + \dots + a_0$  and E(x).

Message "is" points on P(x). (degree n-1, n+2k points.) Channel: Send P(i), receive R(i).

Errors are wrong values at  $\leq k$  points.Error:  $P(i) \neq R(i)$ .

Error locator polynomial:  $E(x) = (x - e_1) \cdot (x - e_k) = x^k + b_{k-1}x^{k-1} + \dots + b_0.$ Find:  $Q(x) = E(x)P(x) = a_{n+k-1}x^{n+k-1} + \dots + a_0$  and E(x). Using n + 2k equations: Q(i) = R(i)E(i).

Message "is" points on P(x). (degree n-1, n+2k points.) Channel: Send P(i), receive R(i).

Errors are wrong values at  $\leq k$  points.Error:  $P(i) \neq R(i)$ .

Error locator polynomial:  $E(x) = (x - e_1) \cdot (x - e_k) = x^k + b_{k-1}x^{k-1} + \dots + b_0.$ Find:  $Q(x) = E(x)P(x) = a_{n+k-1}x^{n+k-1} + \dots + a_0$  and E(x). Using n + 2k equations: Q(i) = R(i)E(i). P(x) = Q(x)/E(x).

For all points  $1, \ldots, i, n+2k = m$ ,

 $Q(i) = R(i)E(i) \pmod{p}$ 

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

$$a_{n+k-1}+\ldots a_0 \equiv R(1)(1+b_{k-1}\cdots b_0) \pmod{p}$$

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

$$a_{n+k-1} + \dots a_0 \equiv R(1)(1 + b_{k-1} \dots b_0) \pmod{p}$$
  
$$a_{n+k-1}(2)^{n+k-1} + \dots a_0 \equiv R(2)((2)^k + b_{k-1}(2)^{k-1} \dots b_0) \pmod{p}$$

÷

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

 $\begin{array}{rcl} a_{n+k-1} + \dots a_0 &\equiv & R(1)(1+b_{k-1}\cdots b_0) \pmod{p} \\ a_{n+k-1}(2)^{n+k-1} + \dots a_0 &\equiv & R(2)((2)^k + b_{k-1}(2)^{k-1}\cdots b_0) \pmod{p} \\ &\vdots \\ a_{n+k-1}(m)^{n+k-1} + \dots a_0 &\equiv & R(m)((m)^k + b_{k-1}(m)^{k-1}\cdots b_0) \pmod{p} \end{array}$ 

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

 $a_{n+k-1} + \dots a_0 \equiv R(1)(1 + b_{k-1} \cdots b_0) \pmod{p}$   $a_{n+k-1}(2)^{n+k-1} + \dots a_0 \equiv R(2)((2)^k + b_{k-1}(2)^{k-1} \cdots b_0) \pmod{p}$   $\vdots$   $a_{n+k-1}(m)^{n+k-1} + \dots a_0 \equiv R(m)((m)^k + b_{k-1}(m)^{k-1} \cdots b_0) \pmod{p}$ 

..and n+2k unknown coefficients of Q(x) and E(x)!

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

 $\begin{array}{rcl} a_{n+k-1} + \dots a_0 &\equiv & R(1)(1 + b_{k-1} \cdots b_0) \pmod{p} \\ a_{n+k-1}(2)^{n+k-1} + \dots a_0 &\equiv & R(2)((2)^k + b_{k-1}(2)^{k-1} \cdots b_0) \pmod{p} \\ &\vdots \\ a_{n+k-1}(m)^{n+k-1} + \dots a_0 &\equiv & R(m)((m)^k + b_{k-1}(m)^{k-1} \cdots b_0) \pmod{p} \end{array}$ 

..and n+2k unknown coefficients of Q(x) and E(x)! Solve for coefficients of Q(x) and E(x).

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

 $\begin{array}{rcl} a_{n+k-1} + \dots a_0 &\equiv & R(1)(1 + b_{k-1} \cdots b_0) \pmod{p} \\ a_{n+k-1}(2)^{n+k-1} + \dots a_0 &\equiv & R(2)((2)^k + b_{k-1}(2)^{k-1} \cdots b_0) \pmod{p} \\ &\vdots \\ a_{n+k-1}(m)^{n+k-1} + \dots a_0 &\equiv & R(m)((m)^k + b_{k-1}(m)^{k-1} \cdots b_0) \pmod{p} \end{array}$ 

..and n+2k unknown coefficients of Q(x) and E(x)! Solve for coefficients of Q(x) and E(x).

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

 $\begin{array}{rcl} a_{n+k-1} + \dots a_0 &\equiv & R(1)(1 + b_{k-1} \cdots b_0) \pmod{p} \\ a_{n+k-1}(2)^{n+k-1} + \dots a_0 &\equiv & R(2)((2)^k + b_{k-1}(2)^{k-1} \cdots b_0) \pmod{p} \\ &\vdots \\ a_{n+k-1}(m)^{n+k-1} + \dots a_0 &\equiv & R(m)((m)^k + b_{k-1}(m)^{k-1} \cdots b_0) \pmod{p} \end{array}$ 

..and n+2k unknown coefficients of Q(x) and E(x)! Solve for coefficients of Q(x) and E(x).

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

 $\begin{array}{rcl} a_{n+k-1} + \dots a_0 &\equiv & R(1)(1 + b_{k-1} \cdots b_0) \pmod{p} \\ a_{n+k-1}(2)^{n+k-1} + \dots a_0 &\equiv & R(2)((2)^k + b_{k-1}(2)^{k-1} \cdots b_0) \pmod{p} \\ &\vdots \\ a_{n+k-1}(m)^{n+k-1} + \dots a_0 &\equiv & R(m)((m)^k + b_{k-1}(m)^{k-1} \cdots b_0) \pmod{p} \end{array}$ 

..and n+2k unknown coefficients of Q(x) and E(x)! Solve for coefficients of Q(x) and E(x).

For all points  $1, \ldots, i, n+2k = m$ ,

$$Q(i) = R(i)E(i) \pmod{p}$$

Gives n + 2k linear equations.

 $\begin{array}{rcl} a_{n+k-1} + \dots a_0 &\equiv & R(1)(1 + b_{k-1} \cdots b_0) \pmod{p} \\ a_{n+k-1}(2)^{n+k-1} + \dots a_0 &\equiv & R(2)((2)^k + b_{k-1}(2)^{k-1} \cdots b_0) \pmod{p} \\ &\vdots \\ a_{n+k-1}(m)^{n+k-1} + \dots a_0 &\equiv & R(m)((m)^k + b_{k-1}(m)^{k-1} \cdots b_0) \pmod{p} \end{array}$ 

..and n+2k unknown coefficients of Q(x) and E(x)! Solve for coefficients of Q(x) and E(x).



What are the ideas?

What are the ideas?

(A) Multiply a incorrect equality by zero makes it correct.

What are the ideas?

(A) Multiply a incorrect equality by zero makes it correct.(B) Multiply by non-zero keeps it informative.

What are the ideas?

(A) Multiply a incorrect equality by zero makes it correct.

- (B) Multiply by non-zero keeps it informative.
- (C) A polynomial of degree k, can have exactly k zeros.

What are the ideas?

- (A) Multiply a incorrect equality by zero makes it correct.
- (B) Multiply by non-zero keeps it informative.
- (C) A polynomial of degree k, can have exactly k zeros.
- (D) Multpliying two polynomials gives a polynomial.

What are the ideas?

- (A) Multiply a incorrect equality by zero makes it correct.
- (B) Multiply by non-zero keeps it informative.
- (C) A polynomial of degree k, can have exactly k zeros.
- (D) Multpliying two polynomials gives a polynomial.
- (E) Q(i) = E(i)R(i) is linear in coeffs of Q and E.

What are the ideas?

- (A) Multiply a incorrect equality by zero makes it correct.
- (B) Multiply by non-zero keeps it informative.
- (C) A polynomial of degree k, can have exactly k zeros.
- (D) Multpliying two polynomials gives a polynomial.
- (E) Q(i) = E(i)R(i) is linear in coeffs of Q and E.

(A), (B), (C).

What are the ideas?

- (A) Multiply a incorrect equality by zero makes it correct.
- (B) Multiply by non-zero keeps it informative.
- (C) A polynomial of degree k, can have exactly k zeros.
- (D) Multpliying two polynomials gives a polynomial.
- (E) Q(i) = E(i)R(i) is linear in coeffs of Q and E.

(A), (B), (C).

 $\implies$  error polynomial.

What are the ideas?

- (A) Multiply a incorrect equality by zero makes it correct.
- (B) Multiply by non-zero keeps it informative.
- (C) A polynomial of degree k, can have exactly k zeros.
- (D) Multpliying two polynomials gives a polynomial.
- (E) Q(i) = E(i)R(i) is linear in coeffs of Q and E.

(A), (B), (C).

 $\implies$  error polynomial.

(D) and (E).

What are the ideas?

- (A) Multiply a incorrect equality by zero makes it correct.
- (B) Multiply by non-zero keeps it informative.
- (C) A polynomial of degree k, can have exactly k zeros.
- (D) Multpliying two polynomials gives a polynomial.
- (E) Q(i) = E(i)R(i) is linear in coeffs of Q and E.

(A), (B), (C).

 $\implies$  error polynomial.

(D) and (E). finish up.

A message of length 4, encoded as P(x) w/packets P(1), ... P(8).

A message of length 4, encoded as P(x) w/packets P(1),...P(8). Recieved packets R(1),...R(8).

A message of length 4, encoded as P(x) w/packets P(1),...P(8). Recieved packets R(1),...R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

A message of length 4, encoded as P(x) w/packets P(1),...P(8). Recieved packets R(1),...R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ (B) The degree of P(x)E(x) = 3 + 2 = 5.

A message of length 4, encoded as P(x) w/packets P(1), ... P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ (B) The degree of P(x)E(x) = 3 + 2 = 5. (C) The degree of E(x) is 2.

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

- (A)  $R(1) \neq P(1)$
- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

(A) E(x) = (x-1)(x-2)

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

(A) 
$$E(x) = (x-1)(x-2)$$

(B) The number of coefficients in E(x) is 2.

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

(A) E(x) = (x-1)(x-2)

- (B) The number of coefficients in E(x) is 2.
- (C) The number of unknown coefficients in E(x) is 2.

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

(A) E(x) = (x-1)(x-2)

- (B) The number of coefficients in E(x) is 2.
- (C) The number of unknown coefficients in E(x) is 2.

(D) 
$$E(x) = (x-1)(x-4)$$

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

(A) 
$$E(x) = (x-1)(x-2)$$

(B) The number of coefficients in E(x) is 2.

(C) The number of unknown coefficients in E(x) is 2.

(D) 
$$E(x) = (x-1)(x-4)$$

(E) 
$$R(4) \neq P(4)$$

A message of length 4, encoded as P(x) w/packets P(1),...P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

(A) 
$$E(x) = (x-1)(x-2)$$

(B) The number of coefficients in E(x) is 2.

(C) The number of unknown coefficients in E(x) is 2.

(D) 
$$E(x) = (x-1)(x-4)$$

(E) 
$$R(4) \neq P(4)$$

(F) The degree of R(x) is 5.

A message of length 4, encoded as P(x) w/packets P(1), ... P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

(A) 
$$E(x) = (x-1)(x-2)$$

(B) The number of coefficients in E(x) is 2.

(C) The number of unknown coefficients in E(x) is 2.

(D) 
$$E(x) = (x-1)(x-4)$$

(E) 
$$R(4) \neq P(4)$$

(F) The degree of R(x) is 5.

(C),(D), (E). (F) doesn't type check!

A message of length 4, encoded as P(x) w/packets P(1), ... P(8).

Recieved packets R(1), ..., R(8).

Packets 1 and 4 are corrupted.

(A)  $R(1) \neq P(1)$ 

- (B) The degree of P(x)E(x) = 3 + 2 = 5.
- (C) The degree of E(x) is 2.
- (D) The number of coefficients of P(x) is 4.
- (E) The number of coefficients of P(x)Q(x) is 6.

(E) is false.

(A) E(x) = (x-1)(x-2)

(B) The number of coefficients in E(x) is 2.

(C) The number of unknown coefficients in E(x) is 2.

(D) 
$$E(x) = (x-1)(x-4)$$

(E) 
$$R(4) \neq P(4)$$

(F) The degree of R(x) is 5.

(C),(D), (E).

(F) doesn't type check! R(i) is the value one recieves.

Communicate *n* packets, with *k* erasures.

How many packets?

Communicate *n* packets, with *k* erasures.

How many packets? n+k

Communicate *n* packets, with *k* erasures.

How many packets? n + kHow to encode?

Communicate *n* packets, with *k* erasures.

How many packets? n+kHow to encode? With polynomial, P(x).

Communicate *n* packets, with *k* erasures.

How many packets? n+kHow to encode? With polynomial, P(x). Of degree?

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
```

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover?
```

Communicate *n* packets, with *k* erasures.

How many packets? n+kHow to encode? With polynomial, P(x). Of degree? n-1Recover? Reconstruct P(x) with any *n* points!

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

How many packets?

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

How many packets? n+2k

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

```
How many packets? n+2k
Why?
```

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

```
How many packets? n+2k
Why?
k extra "good" packets after k errors.
```

Communicate n packets, with k erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

```
How many packets? n+2k
Why?
k extra "good" packets after k errors.
How to encode?
```

Communicate n packets, with k erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

```
How many packets? n+2k
Why?
k extra "good" packets after k errors.
How to encode? With polynomial, P(x).
```

Communicate n packets, with k erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

```
How many packets? n+2k
Why?
k extra "good" packets after k errors.
How to encode? With polynomial, P(x). Of degree?
```

Communicate n packets, with k erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

```
How many packets? n+2k
Why?
k extra "good" packets after k errors.
How to encode? With polynomial, P(x). Of degree? n-1.
```

Communicate n packets, with k erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

```
How many packets? n+2k
Why?
k extra "good" packets after k errors.
How to encode? With polynomial, P(x). Of degree? n-1.
\leq k changes ensure
```

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial.

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Nonlinear equations.

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Nonlinear equations.

Reconstruct E(x) and Q(x) = E(x)P(x).

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Nonlinear equations.

Reconstruct E(x) and Q(x) = E(x)P(x). Linear Equations.

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Nonlinear equations.

Reconstruct E(x) and Q(x) = E(x)P(x). Linear Equations. Polynomial division!

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Nonlinear equations.

Reconstruct E(x) and Q(x) = E(x)P(x). Linear Equations. Polynomial division! P(x) = Q(x)/E(x)!

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Nonlinear equations.

Reconstruct E(x) and Q(x) = E(x)P(x). Linear Equations. Polynomial division! P(x) = Q(x)/E(x)!

Reed-Solomon codes.

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Nonlinear equations.

Reconstruct E(x) and Q(x) = E(x)P(x). Linear Equations. Polynomial division! P(x) = Q(x)/E(x)!

Reed-Solomon codes. Welsh-Berlekamp Decoding.

Communicate *n* packets, with *k* erasures.

```
How many packets? n+k
How to encode? With polynomial, P(x).
Of degree? n-1
Recover? Reconstruct P(x) with any n points!
```

Communicate *n* packets, with *k* errors.

```
How many packets? n+2k
```

Why?

k extra "good" packets after k errors.

How to encode? With polynomial, P(x). Of degree? n-1.

 $\leq k$  changes ensure

two polynomials "correct on" n + k overlap on n points

Two polynomials of degree n-1 mean same polynonmial. Recover?

Reconstruct error polynomial, E(X), and P(x)!

Nonlinear equations.

Reconstruct E(x) and Q(x) = E(x)P(x). Linear Equations. Polynomial division! P(x) = Q(x)/E(x)!

Reed-Solomon codes. Welsh-Berlekamp Decoding. Perfection!



Really Cool!

What's to come?

What's to come? Probability.

What's to come? Probability.

A bag contains:

What's to come? Probability.

A bag contains:



#### What's to come? Probability.

A bag contains:



What is the chance that a ball taken from the bag is blue?

#### What's to come? Probability.

A bag contains:



What is the chance that a ball taken from the bag is blue? Count blue.

#### What's to come? Probability.

A bag contains:



What is the chance that a ball taken from the bag is blue? Count blue. Count total.

#### What's to come? Probability.

A bag contains:



What is the chance that a ball taken from the bag is blue? Count blue. Count total. Divide.

#### What's to come? Probability.

A bag contains:



What is the chance that a ball taken from the bag is blue? Count blue. Count total. Divide.

For now:

#### What's to come? Probability.

A bag contains:



What is the chance that a ball taken from the bag is blue? Count blue. Count total. Divide.

For now: Counting!

#### What's to come? Probability.

A bag contains:



What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide.

For now: Counting!

Later: Probability.

What's to come?

What's to come? Probability.

What's to come? Probability. A bag contains:

What's to come? Probability.

A bag contains:



What's to come? Probability. A bag contains:

What is the chance that a ball taken from the bag is blue?

What's to come? Probability. A bag contains:

What is the chance that a ball taken from the bag is blue? Count blue.

What's to come? Probability. A bag contains:

What is the chance that a ball taken from the bag is blue? Count blue. Count total.

What's to come? Probability. A bag contains:

What is the chance that a ball taken from the bag is blue? Count blue. Count total. Divide. **Chances?** 

What's to come? Probability. A bag contains:

What is the chance that a ball taken from the bag is blue? Count blue. Count total. Divide. **Chances?** (A) Red Probability is 3/8

What's to c	come?	Proba	ability.		
A bag co	ontains	:			
	•				•

What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide. Chances?

(A) Red Probability is 3/8

(B) Blue probability is 3/9

What's to c	come?	Proba	ability.		
A bag co	ntains	:			
					•

What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide. Chances?

(A) Red Probability is 3/8

(B) Blue probability is 3/9

(C) Yellow Probability is 2/8

What's to c	come?	Proba	ability.		
A bag co	ontains	:			
•	•				•

What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide. Chances?

- (A) Red Probability is 3/8
- (B) Blue probability is 3/9
- (C) Yellow Probability is 2/8
- (D) Blue probability is 3/8

What's to c	come?	Proba	ability.		
A bag co	ontains	:			
•	•				•

What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide. Chances?

- (A) Red Probability is 3/8
- (B) Blue probability is 3/9
- (C) Yellow Probability is 2/8
- (D) Blue probability is 3/8

What's to c	come?	Proba	ability.		
A bag co	ontains	:			
	•				•

What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide. Chances?

(A) Red Probability is 3/8

(B) Blue probability is 3/9

(C) Yellow Probability is 2/8

(D) Blue probability is 3/8

Today:

What's to c	come?	Proba	ability.		
A bag co	ontains	:			
	•				•

What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide. Chances?

(A) Red Probability is 3/8

(B) Blue probability is 3/9

- (C) Yellow Probability is 2/8
- (D) Blue probability is 3/8

Today: Counting!

What's to c	come?	Proba	ability.		
A bag co	ontains	:			
•					•

What is the chance that a ball taken from the bag is blue?

Count blue. Count total. Divide. Chances?

(A) Red Probability is 3/8

(B) Blue probability is 3/9

- (C) Yellow Probability is 2/8
- (D) Blue probability is 3/8

Today: Counting!

Combinatorics.

## **Outline: basics**

- 1. Counting.
- 2. Tree
- 3. Rules of Counting
- 4. Sample with/without replacement where order does/doesn't matter.

# Count?


How many outcomes possible for *k* coin tosses? How many poker hands? How many handshakes for *n* people? How many diagonals in a *n* sided convex polygon? How many 10 digit numbers? How many 10 digit numbers without repetition? How many ways can I divide up 5 dollars among 3 people?


How many 3-bit strings?


How many 3-bit strings?


How many different sequences of three bits from  $\{0,1\}$ ?


How many 3-bit strings? How many different sequences of three bits from  $\{0,1\}$ ? How would you make one sequence?
















How many 3-bit strings? How many different sequences of three bits from  $\{0,1\}$ ? How would you make one sequence? How many different ways to do that making?




8 leaves which is  $2 \times 2 \times 2$ . One leaf for each string. 8 3-bit strings! Also recurrence:  $B(n) = 2 \times B(n-1)$ .


How many 3-bit strings? How many different sequences of three bits from  $\{0,1\}$ ? How would you make one sequence? How many different ways to do that making?




8 leaves which is  $2 \times 2 \times 2$ . One leaf for each string. 8 3-bit strings! Also recurrence:  $B(n) = 2 \times B(n-1)$ . B(0) = 1.




 $n_1$ 






Objects made by choosing from  $n_1$ , then  $n_2$ , ..., then  $n_k$  the number of objects is  $n_1 \times n_2 \cdots \times n_k$ .




In picture,  $2 \times 2 \times 3 = 12!$ 

Objects made by choosing from  $n_1$ , then  $n_2$ , ..., then  $n_k$  the number of objects is  $n_1 \times n_2 \cdots \times n_k$ .



In picture,  $2 \times 2 \times 3 = 12!$ Also recurrence:  $S(i) = n_i \times S(i-1)$ ,

Objects made by choosing from  $n_1$ , then  $n_2$ , ..., then  $n_k$  the number of objects is  $n_1 \times n_2 \cdots \times n_k$ .



In picture,  $2 \times 2 \times 3 = 12!$ Also recurrence:  $S(i) = n_i \times S(i-1)$ , Base: S(0) = 1



What's correct?

(A)  $|10 \text{ digit numbers}| = 10^{10}$ 

What's correct? (A) |10 digit numbers| =  $10^{10}$ (B) |k coin tosses| =  $2^{k}$ 

- (A)  $|10 \text{ digit numbers}| = 10^{10}$ (B)  $|k \text{ coin tosses}| = 2^k$
- (C)  $|10 \text{ digit numbers}| = 9 * 10^9$

- (A)  $|10 \text{ digit numbers}| = 10^{10}$ (B)  $|k \text{ coin tosses}| = 2^k$
- (C)  $|10 \text{ digit numbers}| = 9 * 10^9$
- (D)  $|n \text{ digit base } m \text{ numbers}| = m^n$

- (A)  $|10 \text{ digit numbers}| = 10^{10}$
- (B)  $|k \text{ coin tosses}| = 2^k$
- (C)  $|10 \text{ digit numbers}| = 9 * 10^9$
- (D)  $|n \text{ digit base } m \text{ numbers}| = m^n$
- (E)  $|n \text{ digit base } m \text{ numbers}| = (m-1)m^{n-1}$

- (A)  $|10 \text{ digit numbers}| = 10^{10}$
- (B)  $|k \text{ coin tosses}| = 2^k$
- (C)  $|10 \text{ digit numbers}| = 9 * 10^9$
- (D)  $|n \text{ digit base } m \text{ numbers}| = m^n$
- (E)  $|n \text{ digit base } m \text{ numbers}| = (m-1)m^{n-1}$

What's correct?

- (A)  $|10 \text{ digit numbers}| = 10^{10}$
- (B)  $|k \text{ coin tosses}| = 2^k$
- (C)  $|10 \text{ digit numbers}| = 9 * 10^9$
- (D)  $|n \text{ digit base } m \text{ numbers}| = m^n$
- (E) |n digit base *m* numbers  $| = (m-1)m^{n-1}$

(A) or (C)? (D) or (E)? (B) are correct.

How many outcomes possible for k coin tosses?

How many outcomes possible for k coin tosses?

2 ways for first choice,

How many outcomes possible for *k* coin tosses? 2 ways for first choice, 2 ways for second choice, ...

How many outcomes possible for *k* coin tosses? 2 ways for first choice, 2 ways for second choice, ... 2

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2\times 2$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots$ 

How many outcomes possible for *k* coin tosses? 2 ways for first choice, 2 ways for second choice, ...

 $2 \times 2 \cdots \times 2$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice,

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ... 10

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ... 10  $\times$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10\times10\cdots$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10 \times 10 \cdots \times 10$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10 \times 10 \cdots \times 10 = 10^k$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10 \times 10 \cdots \times 10 = 10^k$ 

How many *n* digit base *m* numbers?

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10 \times 10 \cdots \times 10 = 10^k$ 

How many *n* digit base *m* numbers?

m ways for first,

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10 \times 10 \cdots \times 10 = 10^k$ 

How many *n* digit base *m* numbers?

m ways for first, m ways for second, ...

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10 \times 10 \cdots \times 10 = 10^k$ 

How many *n* digit base *m* numbers?

m ways for first, m ways for second, ...  $m^n$ 

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10 \times 10 \cdots \times 10 = 10^k$ 

How many *n* digit base *m* numbers?

m ways for first, m ways for second, ...  $m^n$ 

(Is 09, a two digit number?)

How many outcomes possible for k coin tosses?

2 ways for first choice, 2 ways for second choice, ...  $2 \times 2 \cdots \times 2 = 2^k$ 

How many 10 digit numbers?

10 ways for first choice, 10 ways for second choice, ...  $10 \times 10 \cdots \times 10 = 10^k$ 

How many *n* digit base *m* numbers?

m ways for first, m ways for second, ...  $m^n$ 

(Is 09, a two digit number?)

```
If no. Then (m-1)m^{n-1}.
```

How many functions f mapping S to T?

How many functions f mapping S to T?

|T| ways to choose for  $f(s_1)$ ,

How many functions *f* mapping *S* to *T*?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...

How many functions *f* mapping *S* to *T*?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many functions *f* mapping *S* to *T*?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many polynomials of degree d modulo p?

How many functions *f* mapping *S* to *T*?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many polynomials of degree d modulo p?

p ways to choose for first coefficient,

How many functions *f* mapping *S* to *T*?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many polynomials of degree *d* modulo *p*?

p ways to choose for first coefficient, p ways for second, ...

How many functions *f* mapping *S* to *T*?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many polynomials of degree *d* modulo *p*?

p ways to choose for first coefficient, p ways for second, ...  $...p^{d+1}$ 

How many functions f mapping S to T?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many polynomials of degree d modulo p?

p ways to choose for first coefficient, p ways for second, ...  $...p^{d+1}$ 

p values for first point,

How many functions f mapping S to T?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many polynomials of degree d modulo p?

p ways to choose for first coefficient, p ways for second, ...  $...p^{d+1}$ 

*p* values for first point, *p* values for second, ...

How many functions f mapping S to T?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many polynomials of degree d modulo p?

p ways to choose for first coefficient, p ways for second, ...  $...p^{d+1}$ 

p values for first point, p values for second, ...  $...p^{d+1}$ 

How many functions *f* mapping *S* to *T*?

|T| ways to choose for  $f(s_1)$ , |T| ways to choose for  $f(s_2)$ , ...  $\dots |T|^{|S|}$ 

How many polynomials of degree d modulo p?

p ways to choose for first coefficient, p ways for second, ...  $...p^{d+1}$ 

p values for first point, p values for second, ...  $\dots p^{d+1}$ 

Questions?

<sup>1</sup>By definition: 0! = 1.

How many 10 digit numbers without repeating a digit?

<sup>1</sup>By definition: 0! = 1.

How many 10 digit numbers **without repeating a digit**? 10 ways for first,

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second,

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third,

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size *k* from *n* numbers **without replacement**. *n* ways for first choice,

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size *k* from *n* numbers without replacement.

*n* ways for first choice, n-1 ways for second,

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

```
n ways for first choice, n-1 ways for second, n-2 choices for third,
```

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

How many 10 digit numbers **without repeating a digit**? 10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

... 
$$n * (n-1) * (n-2) \cdot * (n-k+1) = \frac{n!}{(n-k)!}$$
.

How many 10 digit numbers without repeating a digit?

10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

... 
$$n * (n-1) * (n-2) \cdot * (n-k+1) = \frac{n!}{(n-k)!}$$
.

How many orderings of n objects?

How many 10 digit numbers without repeating a digit?

10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

... 
$$n * (n-1) * (n-2) \cdot * (n-k+1) = \frac{n!}{(n-k)!}$$
.

How many orderings of *n* objects?

Permutations of *n* objects.

<sup>&</sup>lt;sup>1</sup>By definition: 0! = 1.

How many 10 digit numbers without repeating a digit?

10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

... 
$$n * (n-1) * (n-2) \cdot * (n-k+1) = \frac{n!}{(n-k)!}$$
.

How many orderings of *n* objects?

#### Permutations of *n* objects.

n ways for first,

<sup>&</sup>lt;sup>1</sup>By definition: 0! = 1.

How many 10 digit numbers without repeating a digit?

10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

... 
$$n * (n-1) * (n-2) \cdot * (n-k+1) = \frac{n!}{(n-k)!}$$
.

How many orderings of n objects?

#### Permutations of *n* objects.

*n* ways for first, n-1 ways for second,

<sup>&</sup>lt;sup>1</sup>By definition: 0! = 1.

How many 10 digit numbers without repeating a digit?

10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

... 
$$n * (n-1) * (n-2) \cdot * (n-k+1) = \frac{n!}{(n-k)!}$$
.

How many orderings of n objects?

#### Permutations of *n* objects.

*n* ways for first, n-1 ways for second, n-2 ways for third,

<sup>&</sup>lt;sup>1</sup>By definition: 0! = 1.

How many 10 digit numbers without repeating a digit?

10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

... 
$$n * (n-1) * (n-2) \cdot * (n-k+1) = \frac{n!}{(n-k)!}$$
.

How many orderings of n objects?

#### Permutations of *n* objects.

*n* ways for first, n-1 ways for second, n-2 ways for third, ...

#### <sup>1</sup>By definition: 0! = 1.

How many 10 digit numbers without repeating a digit?

10 ways for first, 9 ways for second, 8 ways for third, ...

...  $10 * 9 * 8 \cdots * 1 = 10!.^{1}$ 

Sample of size k from n numbers without replacement.

*n* ways for first choice, n-1 ways for second, n-2 choices for third, ...

... 
$$n * (n-1) * (n-2) \cdot * (n-k+1) = \frac{n!}{(n-k)!}$$
.

How many orderings of n objects?

#### Permutations of *n* objects.

*n* ways for first, n-1 ways for second, n-2 ways for third, ...

... 
$$n * (n-1) * (n-2) \cdot *1 = n!$$
.

#### <sup>1</sup>By definition: 0! = 1.

How many one-to-one functions from |S| to |S|.

How many one-to-one functions from |S| to |S|. |S| choices for  $f(s_1)$ ,

How many one-to-one functions from |S| to |S|. |S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

How many one-to-one functions from |S| to |S|. |S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

How many one-to-one functions from |S| to |S|. |S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ... So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ 

How many one-to-one functions from |S| to |S|.

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ A one-to-one function is a permutation!

How many one-to-one functions from |S| to |S|.

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ A one-to-one function is a permutation!

Bijections are permutations.

How many one-to-one functions from |S| to |S|.

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ A one-to-one function is a permutation!

Bijections are permutations.

 $f(x) = 2x \pmod{5}$ 

How many one-to-one functions from |S| to |S|.

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ A one-to-one function is a permutation!

Bijections are permutations.

 $f(x) = 2x \pmod{5}$  $\{0, 1, 2, 3, 4\}$ 

How many one-to-one functions from |S| to |S|.

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ A one-to-one function is a permutation!

Bijections are permutations.

 $f(x) = 2x \pmod{5}$  $\{0, 1, 2, 3, 4\} \rightarrow \{0, 2, 4, 1, 3\}.$ 

How many one-to-one functions from |S| to |S|.

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ A one-to-one function is a permutation!

Bijections are permutations.

 $f(x) = 2x \pmod{5}$  $\{0, 1, 2, 3, 4\} \rightarrow \{0, 2, 4, 1, 3\}.$ 

Switching around elements.

How many one-to-one functions from |S| to |S|.

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ A one-to-one function is a permutation!

Bijections are permutations.

 $f(x) = 2x \pmod{5}$  $\{0, 1, 2, 3, 4\} \rightarrow \{0, 2, 4, 1, 3\}.$ 

Switching around elements.

Number of functions from  $\{0, 1, 2, 3, 4\}$  to  $\{0, 1, 2, 3, 4\}$ ?

How many one-to-one functions from |S| to |S|.

|S| choices for  $f(s_1)$ , |S| - 1 choices for  $f(s_2)$ , ...

So total number is  $|S| \times |S| - 1 \cdots 1 = |S|!$ A one-to-one function is a permutation!

Bijections are permutations.

 $f(x) = 2x \pmod{5}$  $\{0, 1, 2, 3, 4\} \rightarrow \{0, 2, 4, 1, 3\}.$ 

Switching around elements.

Number of functions from  $\{0, 1, 2, 3, 4\}$  to  $\{0, 1, 2, 3, 4\}$ ? 5!.

How many poker hands?

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

How many poker hands?

 $52\times51\times50\times49\times48$ 

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

How many poker hands?

 $52\times51\times50\times49\times48$  ???

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

How many poker hands?

 $52 \times 51 \times 50 \times 49 \times 48$  ???

Are A, K, Q, 10, J of spades and 10, J, Q, K, A of spades the same?

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

How many poker hands?

 $52\times51\times50\times49\times48$  ???

Are A, K, Q, 10, J of spades and 10, J, Q, K, A of spades the same? **Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.<sup>2</sup>

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

How many poker hands?

 $52\times51\times50\times49\times48$  ???

Are A, K, Q, 10, J of spades and 10, J, Q, K, A of spades the same? **Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.<sup>2</sup>

Number of orderings for a poker hand: "5!"

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

How many poker hands?

 $52\times51\times50\times49\times48$  ???

Are A, K, Q, 10, J of spades and 10, J, Q, K, A of spades the same? **Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.<sup>2</sup>

Number of orderings for a poker hand: "5!" (The "!" means factorial, not Exclamation.)

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

How many poker hands?

 $52\times51\times50\times49\times48$  ???

Are A, K, Q, 10, J of spades and 10, J, Q, K, A of spades the same? **Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.<sup>2</sup>

Number of orderings for a poker hand: "5!"

 $\frac{52 \times 51 \times 50 \times 49 \times 48}{5!}$ 

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

# Counting sets..when order doesn't matter.

How many poker hands?

 $52\times51\times50\times49\times48$  ???

Are A, K, Q, 10, J of spades and 10, J, Q, K, A of spades the same? **Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.<sup>2</sup>

Number of orderings for a poker hand: "5!"

Can write as	$52\times51\times50\times49\times48$
	5!
	52!
	$\overline{5! \times 47!}$

<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

# Counting sets..when order doesn't matter.

How many poker hands?

 $52\times51\times50\times49\times48$  ???

Are A, K, Q, 10, J of spades and 10, J, Q, K, A of spades the same? **Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.<sup>2</sup>

Number of orderings for a poker hand: "5!"


Can write as	$52 \times 51 \times 50 \times 49 \times 48$
	5!
	52!
	$\overline{5! \times 47!}$

#### Generic: ways to choose 5 out of 52 possibilities.


<sup>&</sup>lt;sup>2</sup>When each unordered object corresponds equal numbers of ordered objects.

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.




**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.




How many red nodes (ordered objects)?

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.


**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node?


**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.




How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.




How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?  $\frac{9}{3}$ 

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.




How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.




How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

How many poker deals?

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.




How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

How many poker deals?  $52 \cdot 51 \cdot 50 \cdot 49 \cdot 48$ .

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.


How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

How many poker deals?  $52 \cdot 51 \cdot 50 \cdot 49 \cdot 48$ .

How many poker deals per hand?

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.


How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

How many poker deals?  $52 \cdot 51 \cdot 50 \cdot 49 \cdot 48$ .

How many poker deals per hand? Map each deal to ordered deal:

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.


How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

How many poker deals?  $52 \cdot 51 \cdot 50 \cdot 49 \cdot 48$ .

How many poker deals per hand? Map each deal to ordered deal: 5!

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.


How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

How many poker deals?  $52 \cdot 51 \cdot 50 \cdot 49 \cdot 48$ .

How many poker deals per hand? Map each deal to ordered deal: 5!

How many poker hands?

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.


How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

How many poker deals?  $52 \cdot 51 \cdot 50 \cdot 49 \cdot 48$ .

How many poker deals per hand? Map each deal to ordered deal: 5!

How many poker hands?  $\frac{52 \cdot 51 \cdot 50 \cdot 49 \cdot 49}{5!}$ 

**Second Rule of Counting:** If order doesn't matter count ordered objects and then divide by number of orderings.



How many red nodes (ordered objects)? 9.

How many red nodes mapped to one blue node? 3.

How many blue nodes (unordered objects)?  $\frac{9}{3} = 3$ .

How many poker deals?  $52 \cdot 51 \cdot 50 \cdot 49 \cdot 48$ .

How many poker deals per hand? Map each deal to ordered deal: 5! How many poker hands?  $\frac{52 \cdot 51 \cdot 50 \cdot 49 \cdot 48}{5!}$ Questions?

$$n \times (n-1)$$

$$\frac{n \times (n-1)}{2}$$

$$\frac{n\times(n-1)}{2}=\frac{n!}{(n-2)!\times 2}$$

Choose 2 out of n?

$$\frac{n\times(n-1)}{2}=\frac{n!}{(n-2)!\times 2}$$

Choose 2 out of n?

$$\frac{n\times(n-1)}{2}=\frac{n!}{(n-2)!\times 2}$$

$$n \times (n-1) \times (n-2)$$

Choose 2 out of n?

$$\frac{n\times(n-1)}{2}=\frac{n!}{(n-2)!\times 2}$$

$$\frac{n\times(n-1)\times(n-2)}{3!}$$

Choose 2 out of n?

$$\frac{n\times(n-1)}{2}=\frac{n!}{(n-2)!\times 2}$$

$$\frac{n \times (n-1) \times (n-2)}{3!} = \frac{n!}{(n-3)! \times 3!}$$

Choose 2 out of n?

$$\frac{n\times(n-1)}{2}=\frac{n!}{(n-2)!\times 2}$$

Choose 3 out of n?

$$\frac{n \times (n-1) \times (n-2)}{3!} = \frac{n!}{(n-3)! \times 3!}$$

Choose k out of n?

 $\frac{n!}{(n-k)!}$ 

Choose 2 out of n?

$$\frac{n\times(n-1)}{2}=\frac{n!}{(n-2)!\times 2}$$

Choose 3 out of n?

$$\frac{n \times (n-1) \times (n-2)}{3!} = \frac{n!}{(n-3)! \times 3!}$$

Choose k out of n?

 $\frac{n!}{(n-k)!}$ 

Choose 2 out of n?

$$\frac{n\times(n-1)}{2}=\frac{n!}{(n-2)!\times 2}$$

Choose 3 out of n?

$$\frac{n \times (n-1) \times (n-2)}{3!} = \frac{n!}{(n-3)! \times 3!}$$

Choose k out of n?

 $\frac{n!}{(n-k)! \times k!}$ 

Choose 2 out of n?

$$\frac{n\times(n-1)}{2} = \frac{n!}{(n-2)!\times 2}$$

Choose 3 out of n?

$$\frac{n \times (n-1) \times (n-2)}{3!} = \frac{n!}{(n-3)! \times 3!}$$

Choose k out of n?

 $\frac{n!}{(n-k)! \times k!}$ 

Notation:  $\binom{n}{k}$  and pronounced "*n* choose *k*."

Choose 2 out of n?

$$\frac{n\times(n-1)}{2} = \frac{n!}{(n-2)!\times 2}$$

Choose 3 out of n?

$$\frac{n \times (n-1) \times (n-2)}{3!} = \frac{n!}{(n-3)! \times 3!}$$

Choose k out of n?

 $\frac{n!}{(n-k)! \times k!}$ 

Notation:  $\binom{n}{k}$  and pronounced "*n* choose *k*." Familiar?

Choose 2 out of n?

$$\frac{n\times(n-1)}{2} = \frac{n!}{(n-2)!\times 2}$$

Choose 3 out of n?

$$\frac{n \times (n-1) \times (n-2)}{3!} = \frac{n!}{(n-3)! \times 3!}$$


Choose k out of n?

 $\frac{n!}{(n-k)! \times k!}$ 

Notation:  $\binom{n}{k}$  and pronounced "*n* choose *k*." Familiar? Questions?


### Example: Visualize the proof..

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




#### Example: Visualize the proof..

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals: 52

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51$ 

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52\times51\times50$ 

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ .

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule.

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ? Hand: Q, K, A.

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ? Hand: Q, K, A. Deals: Q, K, A:

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ? Hand: Q, K, A. Deals: Q, K, A : Q, A, K:

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ? Hand: Q, K, A. Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K.

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ? Hand: Q, K, A. Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K.  $\Delta = 3 \times 2 \times 1$ 

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ? Hand: Q, K, A.

Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K.

 $\Delta = 3 \times 2 \times 1$  First rule again.

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: Q, K, A. Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K.  $\Delta = 3 \times 2 \times 1$  First rule again.

Total:

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: Q, K, A. Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K.

 $\Delta = 3 \times 2 \times 1$  First rule again. Total:  $\frac{52!}{49!3!}$ 

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?


Hand: *Q*,*K*,*A*.

Deals: *Q*,*K*,*A* : *Q*,*A*,*K* : *K*,*A*,*Q* : *K*,*A*,*Q* : *A*,*K*,*Q* : *A*,*Q*,*K*.

 $\Delta = 3 \times 2 \times 1$  First rule again.

Total:  $\frac{52!}{49!3!}$  Second Rule!

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: Q, K, A. Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K.  $\Delta = 3 \times 2 \times 1$  First rule again. Total:  $\frac{52!}{40!3!}$  Second Rule!

Choose k out of n.

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: *Q*,*K*,*A*.

Deals: *Q*,*K*,*A* : *Q*,*A*,*K* : *K*,*A*,*Q* : *K*,*A*,*Q* : *A*,*K*,*Q* : *A*,*Q*,*K*.


 $\Delta = 3 \times 2 \times 1$  First rule again.

Total:  $\frac{52!}{49|3|}$  Second Rule!

Choose k out of n.

Ordered set:  $\frac{n!}{(n-k)!}$ 

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: *Q*,*K*,*A*.

Deals: *Q*, *K*, *A* : *Q*, *A*, *K* : *K*, *A*, *Q* : *K*, *A*, *Q* : *A*, *K*, *Q* : *A*, *Q*, *K*.


 $\Delta = 3 \times 2 \times 1$  First rule again.

Total:  $\frac{52!}{49!3!}$  Second Rule!

Choose k out of n.

Ordered set:  $\frac{n!}{(n-k)!}$  Orderings of one hand?

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: *Q*,*K*,*A*.

Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K.


 $\Delta = 3 \times 2 \times 1$  First rule again.

Total:  $\frac{52!}{49!3!}$  Second Rule!

Choose k out of n.

Ordered set:  $\frac{n!}{(n-k)!}$  Orderings of one hand? k!

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: Q,K,A.

Deals: Q, K, A : Q, A, K : K, A, Q : K, A, Q : A, K, Q : A, Q, K.


 $\Delta = 3 \times 2 \times 1$  First rule again.

Total:  $\frac{52!}{49!3!}$  Second Rule!

Choose k out of n.

Ordered set:  $\frac{n!}{(n-k)!}$  Orderings of one hand? k! (By first rule!)

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: *Q*,*K*,*A*.

Deals: *Q*, *K*, *A* : *Q*, *A*, *K* : *K*, *A*, *Q* : *K*, *A*, *Q* : *A*, *K*, *Q* : *A*, *Q*, *K*.


 $\Delta = 3 \times 2 \times 1$  First rule again.

Total: <sup>52!</sup>/<sub>49!3!</sub> Second Rule!

Choose k out of n.

Ordered set:  $\frac{n!}{(n-k)!}$  Orderings of one hand? *k*! (By first rule!)  $\implies$  Total:  $\frac{n!}{(n-k)!k!}$ 

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: *Q*,*K*,*A*.

Deals: *Q*, *K*, *A* : *Q*, *A*, *K* : *K*, *A*, *Q* : *K*, *A*, *Q* : *A*, *K*, *Q* : *A*, *Q*, *K*.


 $\Delta = 3 \times 2 \times 1$  First rule again.

Total: <sup>52!</sup>/<sub>49!3!</sub> Second Rule!

Choose k out of n.

Ordered set:  $\frac{n!}{(n-k)!}$  Orderings of one hand? *k*! (By first rule!)  $\implies$  Total:  $\frac{n!}{(n-k)!k!}$  Second rule.

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



3 card Poker deals:  $52 \times 51 \times 50 = \frac{52!}{49!}$ . First rule. Poker hands:  $\Delta$ ?

Hand: *Q*,*K*,*A*.

Deals: *Q*, *K*, *A* : *Q*, *A*, *K* : *K*, *A*, *Q* : *K*, *A*, *Q* : *A*, *K*, *Q* : *A*, *Q*, *K*.


 $\Delta = 3 \times 2 \times 1$  First rule again.

Total: <sup>52!</sup>/<sub>49!3!</sub> Second Rule!


Choose k out of n.

Ordered set:  $\frac{n!}{(n-k)!}$  Orderings of one hand? *k*! (By first rule!)  $\implies$  Total:  $\frac{n!}{(n-k)!k!}$  Second rule.

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM?

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7!

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule.

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same!

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ?

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM A<sub>1</sub>NA<sub>2</sub>GRA<sub>3</sub>M,

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM A<sub>1</sub>NA<sub>2</sub>GRA<sub>3</sub>M, A<sub>2</sub>NA<sub>1</sub>GRA<sub>3</sub>M,

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM A<sub>1</sub>NA<sub>2</sub>GRA<sub>3</sub>M, A<sub>2</sub>NA<sub>1</sub>GRA<sub>3</sub>M, ....

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM A<sub>1</sub>NA<sub>2</sub>GRA<sub>3</sub>M, A<sub>2</sub>NA<sub>1</sub>GRA<sub>3</sub>M, ...  $\Delta = 3 \times 2 \times 1$ 

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...




Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM A<sub>1</sub>NA<sub>2</sub>GRA<sub>3</sub>M, A<sub>2</sub>NA<sub>1</sub>GRA<sub>3</sub>M, ...  $\Delta = 3 \times 2 \times 1 = 3!$ 

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM A<sub>1</sub>NA<sub>2</sub>GRA<sub>3</sub>M, A<sub>2</sub>NA<sub>1</sub>GRA<sub>3</sub>M, ...  $\Delta = 3 \times 2 \times 1 = 3!$  First rule!


First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM A<sub>1</sub>NA<sub>2</sub>GRA<sub>3</sub>M, A<sub>2</sub>NA<sub>1</sub>GRA<sub>3</sub>M, ...  $\Delta = 3 \times 2 \times 1 = 3!$  First rule!  $\implies \frac{7!}{3!}$ 

#### Example: Anagram

First rule:  $n_1 \times n_2 \cdots \times n_3$ . Product Rule. Second rule: when order doesn't matter divide...



Orderings of ANAGRAM? Ordered Set: 7! First rule. A's are the same! What is  $\Delta$ ? ANAGRAM A<sub>1</sub>NA<sub>2</sub>GRA<sub>3</sub>M, A<sub>2</sub>NA<sub>1</sub>GRA<sub>3</sub>M, ...  $\Delta = 3 \times 2 \times 1 = 3!$  First rule!  $\implies \frac{7!}{3!}$  Second rule!

(A) |Poker hands| =  $\binom{52}{5}$ 

(A) |Poker hands| =  $\binom{52}{5}$ (B) Orderings of ANAGRAM = 7!/3! (A)  $|Poker hands| = {52 \choose 5}$ (B) Orderings of ANAGRAM = 7!/3! (C) Orderings of "CAT" = 3! (A)  $|Poker hands| = {52 \choose 5}$ (B) Orderings of ANAGRAM = 7!/3! (C) Orderings of "CAT" = 3! (D) Orders of MISSISSIPPI = 11!/4!4!2!

(A)  $|Poker hands| = {52 \choose 5}$ (B) Orderings of ANAGRAM = 7!/3! (C) Orderings of "CAT" = 3! (D) Orders of MISSISSIPPI = 11!/4!4!2! (E) Orderings of ANAGRAM = 7!/4!

(A)  $|Poker hands| = {52 \choose 5}$ (B) Orderings of ANAGRAM = 7!/3! (C) Orderings of "CAT" = 3! (D) Orders of MISSISSIPPI = 11!/4!4!2! (E) Orderings of ANAGRAM = 7!/4!

(F) Orders of MISSISSIPPI = 11!/10!

(A)  $|Poker hands| = {52 \choose 5}$ (B) Orderings of ANAGRAM = 7!/3! (C) Orderings of "CAT" = 3! (D) Orders of MISSISSIPPI = 11!/4!4!2! (E) Orderings of ANAGRAM = 7!/4! (F) Orders of MISSISSIPPI = 11!/10!

(A)-(E) are correct.

How many orderings of letters of CAT?

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies 3 \times 2 \times 1$ 

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3 × 2 × 1 = 3! orderings

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3 × 2 × 1 = 3! orderings

How many orderings of the letters in ANAGRAM?

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3 × 2 × 1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered,

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3 × 2 × 1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3 × 2 × 1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters.

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3 × 2 × 1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7!

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7! total "extra counts" or orderings of three A's?

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7! total "extra counts" or orderings of three A's? 3!

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7!

total "extra counts" or orderings of three A's? 3!

Total orderings?

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7! total "extra counts" or orderings of three A's? 3!

Total orderings?  $\frac{7!}{3!}$ 

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7! total "extra counts" or orderings of three A's? 3!

Total orderings?  $\frac{7!}{3!}$ 

How many orderings of MISSISSIPPI?

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7! total "extra counts" or orderings of three A's? 3!

Total orderings?  $\frac{7!}{3!}$ 

How many orderings of MISSISSIPPI?

4 S's, 4 I's, 2 P's.

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7!

total "extra counts" or orderings of three A's? 3!

Total orderings?  $\frac{7!}{3!}$ 

How many orderings of MISSISSIPPI?

4 S's, 4 l's, 2 P's.

11 letters total.

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7!

total "extra counts" or orderings of three A's? 3!

Total orderings?  $\frac{7!}{3!}$ 

How many orderings of MISSISSIPPI?

4 S's, 4 I's, 2 P's.

11 letters total.

11! ordered objects.

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7!

total "extra counts" or orderings of three A's? 3!

Total orderings?  $\frac{7!}{3!}$ 

How many orderings of MISSISSIPPI?

4 S's, 4 I's, 2 P's.

11 letters total.

11! ordered objects.

 $4! \times 4! \times 2!$  ordered objects per "unordered object"

How many orderings of letters of CAT?

3 ways to choose first letter, 2 ways for second, 1 for last.

 $\implies$  3  $\times$  2  $\times$  1 = 3! orderings

How many orderings of the letters in ANAGRAM?

Ordered, except for A!

total orderings of 7 letters. 7!

total "extra counts" or orderings of three A's? 3!

Total orderings?  $\frac{7!}{3!}$ 

How many orderings of MISSISSIPPI?

4 S's, 4 I's, 2 P's.

11 letters total.

11! ordered objects.

 $4! \times 4! \times 2!$  ordered objects per "unordered object"

 $\implies \frac{11!}{4!4!2!}.$ 

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

k Samples with replacement from *n* items:  $n^k$ .

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

```
k Samples with replacement from n items: n^k.
Sample without replacement: \frac{n!}{(n-k)!}
```

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

Second rule: when order doesn't matter (sometimes) can divide...

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

# Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

# Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

**One-to-one rule: equal in number if one-to-one correspondence.** pause Bijection!

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

# Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

**One-to-one rule: equal in number if one-to-one correspondence.** pause Bijection!

Sample *k* times from *n* objects with replacement and order doesn't matter:  $\binom{k+n-1}{n-1}$ .

## Sampling...

Sample k items out of n

Sample *k* items out of *n* Without replacement:

Sample *k* items out of *n* Without replacement: Order matters:

Sample *k* items out of *n* Without replacement: Order matters:  $n \times$ 

Sample k items out of n

Without replacement: Order matters:  $n \times n - 1 \times n - 2 \dots$ 

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1$ 

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ 

Order does not matter:

Second Rule: divide by number of orders

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ 

Order does not matter:

Second Rule: divide by number of orders – "k!"

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Second Rule: divide by number of orders - "k!"

$$\implies \frac{n!}{(n-k)!k!}.$$

Sample *k* items out of *n* Without replacement: Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter: Second Rule: divide by number of orders - "*k*!"  $\implies \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

Sample k items out of n Without replacement: Order matters:  $n \times n = 1 \times n = 2$ 

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter: Second Rule: divide by number of orders – "*k*!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Sample *k* items out of *n* Without replacement: Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter: Second Rule: divide by number of orders - "*k*!"  $\implies \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

With Replacement. Order matters: *n* 

Sample *k* items out of *n* Without replacement: Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter: Second Rule: divide by number of orders - "*k*!"  $\implies \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

With Replacement. Order matters:  $n \times n$ 

Sample *k* items out of *n* Without replacement: Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter: Second Rule: divide by number of orders - "*k*!"  $\implies \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

With Replacement. Order matters:  $n \times n \times ... n$ 

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times ... n = n^k$ Order does not matter:

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Second Pule: divide by number of

Second Rule: divide by number of orders – "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times ... n = n^k$ Order does not matter: Second rule

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Second Rule: divide by number of orders – "k!"

 $\implies \frac{n!}{(n-k)!k!}$ . "n choose k"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ Order does not matter: Second rule ???

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Second Rule: divide by number of orders – "k!"

 $\implies \frac{n!}{(n-k)!k!}$ . "n choose k"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ Order does not matter: Second rule ???

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

Problem: depends on how many of each item we chose!

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ 

Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ 

Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Second Rule: divide by number of orders – "k!"

 $\implies \frac{n!}{(n-k)!k!}$ "n choose k"

With Replacement.

Order matters:  $n \times n \times ... n = n^k$ 

Order does not matter: Second rule ???

Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Unordered elt: 1,2,3

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ 

Order does not matter:

Second Rule: divide by number of orders – "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Unordered elt: 1,2,3 3! ordered elts map to it.

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ 

Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

#### Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Unordered elt: 1,2,3 3! ordered elts map to it. Unordered elt: 1,2,2

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

# Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Unordered elt: 1,2,33! ordered elts map to it.Unordered elt: 1,2,2 $\frac{3!}{2!}$  ordered elts map to it.

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

# Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Unordered elt: 1,2,33! ordered elts map to it.Unordered elt: 1,2,2 $\frac{3!}{2!}$  ordered elts map to it.

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Order does not matter:

Second Rule: divide by number of orders – "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

# Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Unordered elt: 1,2,3 3! ordered elts map to it.

Unordered elt: 1,2,2  $\frac{3!}{2!}$  ordered elts map to it.

How do we deal with this

Sample k items out of n

Without replacement:

Order matters:  $n \times n - 1 \times n - 2 \dots \times n - k + 1 = \frac{n!}{(n-k)!}$ Order does not matter:

Order does not matter:

Second Rule: divide by number of orders - "k!"

 $\implies \frac{n!}{(n-k)!k!}.$ "*n* choose *k*"

With Replacement.

Order matters:  $n \times n \times \ldots n = n^k$ 

Order does not matter: Second rule ???

## Problem: depends on how many of each item we chose!

Different number of unordered elts map to each unordered elt.

Unordered elt: 1,2,3 3! ordered elts map to it.

Unordered elt: 1,2,2  $\frac{3!}{2!}$  ordered elts map to it.

How do we deal with this mess??

How many ways can Bob and Alice split 5 dollars?

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or  $Alice(2^5)$ , divide out order

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or  $Alice(2^5)$ , divide out order ???

5 dollars for Bob and 0 for Alice:

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

4 for Bob and 1 for Alice:

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

4 for Bob and 1 for Alice:

5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

4 for Bob and 1 for Alice: 5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...

"Sorted" way to specify, first Alice's dollars, then Bob's.

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

4 for Bob and 1 for Alice: 5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...

"Sorted" way to specify, first Alice's dollars, then Bob's. (B, B, B, B, B): (A, B, B, B, B): (A, A, B, B, B):

How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

4 for Bob and 1 for Alice: 5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...

"Sorted" way to specify, first Alice's dollars, then Bob's. (B, B, B, B, B): (A, B, B, B, B): (A, A, B, B, B): (A, A, B, B, B): and so on.


How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

4 for Bob and 1 for Alice: 5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...

"Sorted" way to specify, first Alice's dollars, then Bob's.

(B, B, B, B, B, B): (A, B, B, B, B): (A, A, B, B, B): and so on.




How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

4 for Bob and 1 for Alice: 5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...

"Sorted" way to specify, first Alice's dollars, then Bob's. (B,B,B,B,B): 1: (B,B,B,B,B) (A,B,B,B,B): (A,A,B,B,B): and so on.




How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).

4 for Bob and 1 for Alice: 5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...

"Sorted" way to specify, first Alice's dollars, then Bob's.
(*B*, *B*, *B*, *B*, *B*): 1: (B, B, B, B, B)
(*A*, *B*, *B*, *B*, *B*): 5: (A, B, B, B, B), (B, A, B, B, B), (B, B, A, B, B), ...
(*A*, *A*, *B*, *B*, *B*):

and so on.



How many ways can Bob and Alice split 5 dollars? For each of 5 dollars pick Bob or Alice(2<sup>5</sup>), divide out order ???

5 dollars for Bob and 0 for Alice: one ordered set: (B, B, B, B, B).


4 for Bob and 1 for Alice: 5 ordered sets: (*A*, *B*, *B*, *B*, *B*); (*B*, *A*, *B*, *B*, *B*); ...


"Sorted" way to specify, first Alice's dollars, then Bob's.

(*B*, *B*, *B*, *B*, *B*, *B*): 1: (B, B, B, B, B)

(*A*, *B*, *B*, *B*, *B*): 5: (A,B,B,B,B),(B,A,B,B,B),(B,B,A,B,B),...

(A, A, B, B, B):  $\binom{5}{2}$ ; (A, A, B, B, B), (A, B, A, B, B), (A, B, B, A, B), ... and so on.





Second rule of counting is no good here!

How many ways can Alice, Bob, and Eve split 5 dollars.

How many ways can Alice, Bob, and Eve split 5 dollars. Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).

How many ways can Alice, Bob, and Eve split 5 dollars.

Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).

Separate Alice's dollars from Bob's and then Bob's from Eve's.

How many ways can Alice, Bob, and Eve split 5 dollars.

Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

How many ways can Alice, Bob, and Eve split 5 dollars.

Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

Alice: 2, Bob: 1, Eve: 2.

How many ways can Alice, Bob, and Eve split 5 dollars.

```
Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).
```

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

Alice: 2, Bob: 1, Eve: 2. Stars and Bars:  $\star \star |\star| \star \star$ .

How many ways can Alice, Bob, and Eve split 5 dollars.

```
Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).
```

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

Alice: 2, Bob: 1, Eve: 2.

Stars and Bars:  $\star \star |\star| \star \star$ .

Alice: 0, Bob: 1, Eve: 4.

How many ways can Alice, Bob, and Eve split 5 dollars.

```
Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).
```

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

Alice: 2, Bob: 1, Eve: 2. Stars and Bars:  $\star \star |\star| \star \star$ .

Alice: 0, Bob: 1, Eve: 4. Stars and Bars: |\*|\*\*\*\*.

How many ways can Alice, Bob, and Eve split 5 dollars.

```
Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).
```

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

Alice: 2, Bob: 1, Eve: 2. Stars and Bars: \*\*|\*|\*\*.

Alice: 0, Bob: 1, Eve: 4. Stars and Bars: |\*|\*\*\*\*.

Each split "is" a sequence of stars and bars.

How many ways can Alice, Bob, and Eve split 5 dollars.

```
Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).
```

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

Alice: 2, Bob: 1, Eve: 2. Stars and Bars: \*\*|\*|\*\*.

Alice: 0, Bob: 1, Eve: 4.

Stars and Bars:  $|\star| \star \star \star \star$ .

Each split "is" a sequence of stars and bars. Each sequence of stars and bars "is" a split.

How many ways can Alice, Bob, and Eve split 5 dollars.

```
Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).
```

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

Alice: 2, Bob: 1, Eve: 2. Stars and Bars: \*\*|\*|\*\*.

Alice: 0, Bob: 1, Eve: 4.

Stars and Bars:  $|\star| \star \star \star \star$ .

Each split "is" a sequence of stars and bars. Each sequence of stars and bars "is" a split.

How many ways can Alice, Bob, and Eve split 5 dollars.

```
Alice gets 3, Bob gets 1, Eve gets 1: (A, A, A, B, E).
```

Separate Alice's dollars from Bob's and then Bob's from Eve's.

Five dollars are five stars: \*\*\*\*.

Alice: 2, Bob: 1, Eve: 2. Stars and Bars: \*\*|\*|\*\*.

Alice: 0, Bob: 1, Eve: 4. Stars and Bars: |\*|\*\*\*\*.

Each split "is" a sequence of stars and bars. Each sequence of stars and bars "is" a split.

Counting Rule: if there is a one-to-one mapping between two sets they have the same size!

How many different 5 star and 2 bar diagrams?

How many different 5 star and 2 bar diagrams?

| \* | \* \* \* \*.

How many different 5 star and 2 bar diagrams?

\* \* \* \* \* \*.

7 positions in which to place the 2 bars.

\_\_\_\_\_

How many different 5 star and 2 bar diagrams?

\* \* \* \* \* \*.

\_ \_ \_ \_ \_ \_ \_

7 positions in which to place the 2 bars.

Alice: 0; Bob 1; Eve: 4

How many different 5 star and 2 bar diagrams?

\* \* \* \* \* \*.

\_ \_ \_ \_ \_ \_ \_

7 positions in which to place the 2 bars.

```
Alice: 0; Bob 1; Eve: 4
```

How many different 5 star and 2 bar diagrams?

\* \* \* \* \* \*.

- - - - - - -

7 positions in which to place the 2 bars.

Alice: 0; Bob 1; Eve: 4  $| \star | \star \star \star \star$ . Bars in first and third position.

How many different 5 star and 2 bar diagrams?

\* \* \* \* \* \*.

\_ \_ \_ \_ \_ \_ \_

7 positions in which to place the 2 bars.

Alice: 0; Bob 1; Eve: 4  $| \star | \star \star \star \star$ . Bars in first and third position. Alice: 1; Bob 4; Eve: 0

How many different 5 star and 2 bar diagrams?

\* \* \* \* \* \*.

\_ \_ \_ \_ \_ \_ \_

7 positions in which to place the 2 bars.

```
Alice: 0; Bob 1; Eve: 4
| * | * * * *.
Bars in first and third position.
Alice: 1; Bob 4; Eve: 0
* | * * * * |.
```

How many different 5 star and 2 bar diagrams?

\* \* \* \* \* \*.

\_ \_ \_ \_ \_ \_

7 positions in which to place the 2 bars.

Alice: 0; Bob 1; Eve: 4  $| \star | \star \star \star \star$ . Bars in first and third position. Alice: 1; Bob 4; Eve: 0  $\star | \star \star \star \star |$ . Bars in second and seventh position.

How many different 5 star and 2 bar diagrams?

| \* | \* \* \* \*.

\_ \_ \_ \_ \_ \_

7 positions in which to place the 2 bars.

Alice: 0; Bob 1; Eve: 4  $| \star | \star \star \star \star$ . Bars in first and third position. Alice: 1; Bob 4; Eve: 0  $\star | \star \star \star \star |$ . Bars in second and seventh position.  $\binom{7}{2}$  ways to do so and

How many different 5 star and 2 bar diagrams?

| \* | \* \* \* \*.

\_ \_ \_ \_ \_ \_

7 positions in which to place the 2 bars.

Alice: 0; Bob 1; Eve: 4  $| \star | \star \star \star \star$ . Bars in first and third position.

```
Alice: 1; Bob 4; Eve: 0
```

\* | \* \* \* \* |.

Bars in second and seventh position.

 $\binom{7}{2}$  ways to do so and

 $\binom{7}{2}$  ways to split 5 dollars among 3 people.

Ways to add up *n* numbers to sum to *k*?

Ways to add up *n* numbers to sum to *k*? or

" k from n with replacement where order doesn't matter."

Ways to add up *n* numbers to sum to *k*? or

" *k* from *n* with replacement where order doesn't matter." In general, *k* stars n-1 bars.

\*\* \* ... \*\*.

Ways to add up *n* numbers to sum to *k*? or

" *k* from *n* with replacement where order doesn't matter." In general, *k* stars n-1 bars.

\*\* \* ... \*\*.

n+k-1 positions from which to choose n-1 bar positions.

Ways to add up *n* numbers to sum to *k*? or

" *k* from *n* with replacement where order doesn't matter." In general, *k* stars n-1 bars.

\*\* \* ··· \*\*.

n+k-1 positions from which to choose n-1 bar positions.

 $\binom{n+k-1}{n-1}$ 

Ways to add up *n* numbers to sum to *k*? or

" *k* from *n* with replacement where order doesn't matter." In general, *k* stars n-1 bars.

\*\* \* ··· \*\*.

n+k-1 positions from which to choose n-1 bar positions.

 $\binom{n+k-1}{n-1}$ 

Or: *k* unordered choices from set of *n* possibilities with replacement. **Sample with replacement where order doesn't matter.** 

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

k Samples with replacement from *n* items:  $n^k$ .

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

```
k Samples with replacement from n items: n^k.
Sample without replacement: \frac{n!}{(n-k)!}
```

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

Second rule: when order doesn't matter (sometimes) can divide...

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

# Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

# Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

**One-to-one rule: equal in number if one-to-one correspondence.** pause Bijection!

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

# Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

**One-to-one rule: equal in number if one-to-one correspondence.** pause Bijection!

Sample *k* times from *n* objects with replacement and order doesn't matter:  $\binom{k+n-1}{n-1}$ .



Mark whats correct.

## Poll

#### Mark whats correct.

- (A) ways to split *k* dollars among *n*:  $\binom{k+n-1}{n-1}$
- (B) ways to split *n* dollars among *k*:  $\binom{n+k-1}{k-1}$
- (C) ways to split 5 dollars among 3:  $\binom{5+3-1}{3-1}$

(D) ways to split 5 dollars among  $3:\binom{7}{5}$ 

# Poll

#### Mark whats correct.

- (A) ways to split *k* dollars among *n*:  $\binom{k+n-1}{n-1}$
- (B) ways to split *n* dollars among *k*:  $\binom{n+k-1}{k-1}$
- (C) ways to split 5 dollars among 3:  $\binom{5+3-1}{3-1}$
- (D) ways to split 5 dollars among  $3:\binom{7}{5}$

All correct.

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

k Samples with replacement from *n* items:  $n^k$ .

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

Second rule: when order doesn't matter divide..when possible.

#### **First rule:** $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

#### Second rule: when order doesn't matter divide..when possible.

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

#### **First rule:** $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

#### Second rule: when order doesn't matter divide..when possible.

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

#### One-to-one rule: equal in number if one-to-one correspondence.

#### **First rule:** $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

#### Second rule: when order doesn't matter divide..when possible.

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

#### One-to-one rule: equal in number if one-to-one correspondence.

Sample with replacement and order doesn't matter:  $\binom{k+n-1}{n-1}$ .

#### **First rule:** $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

#### Second rule: when order doesn't matter divide..when possible.

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

#### One-to-one rule: equal in number if one-to-one correspondence.

Sample with replacement and order doesn't matter:  $\binom{k+n-1}{n-1}$ . Distribute *k* samples (stars) over *n* poss. (*n*-1 bars group poss..)

#### **First rule:** $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

#### Second rule: when order doesn't matter divide..when possible.

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

#### One-to-one rule: equal in number if one-to-one correspondence.

Sample with replacement and order doesn't matter:  $\binom{k+n-1}{n-1}$ . Distribute *k* samples (stars) over *n* poss. (*n* – 1 bars group poss..) Distribute *k* dollars to *n* people.

#### **First rule:** $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

#### Second rule: when order doesn't matter divide..when possible.

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

#### One-to-one rule: equal in number if one-to-one correspondence.

Sample with replacement and order doesn't matter:  $\binom{k+n-1}{n-1}$ . Distribute *k* samples (stars) over *n* poss. (*n* – 1 bars group poss..) Distribute *k* dollars to *n* people.

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

k Samples with replacement from *n* items:  $n^k$ .

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

**First rule:**  $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

Second rule:

when order doesn't matter (sometimes) can divide...

#### **First rule:** $n_1 \times n_2 \cdots \times n_3$ .

*k* Samples with replacement from *n* items:  $n^k$ . Sample without replacement:  $\frac{n!}{(n-k)!}$ 

### Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

```
First rule: n_1 \times n_2 \cdots \times n_3.
```

```
k Samples with replacement from n items: n^k.
Sample without replacement: \frac{n!}{(n-k)!}
```

### Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

One-to-one rule: equal in number if one-to-one correspondence.

```
First rule: n_1 \times n_2 \cdots \times n_3.
```

```
k Samples with replacement from n items: n^k.
Sample without replacement: \frac{n!}{(n-k)!}
```

### Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

### One-to-one rule: equal in number if one-to-one correspondence. Bijection!

```
First rule: n_1 \times n_2 \cdots \times n_3.
```

```
k Samples with replacement from n items: n^k.
Sample without replacement: \frac{n!}{(n-k)!}
```

### Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

### One-to-one rule: equal in number if one-to-one correspondence. Bijection!

Sample k n objects with replacement and order doesn't matter:

```
First rule: n_1 \times n_2 \cdots \times n_3.
```

```
k Samples with replacement from n items: n^k.
Sample without replacement: \frac{n!}{(n-k)!}
```

### Second rule: when order doesn't matter (sometimes) can divide...

Sample without replacement and order doesn't matter:  $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ . "*n* choose *k*"

### One-to-one rule: equal in number if one-to-one correspondence. Bijection!

Sample k n objects with replacement and order doesn't matter:

 $\binom{k+n-1}{n-1}$ .