

## Lecture 16: Continuing Probability.

Events, Conditional Probability, Independence, Bayes' Rule

## Probability Basics:Poll

What is a probability space?

- (A) A set and a function on the elements.
  - (B) The values of the function are real numbers.
  - (C) The values of the function are positive integers.
  - (D) An element of the set is an outcome.
  - (E) There is an experiment associated with a probability space.
  - (F) The values in the set are integers.
- (A),(B), (D), (E).

# Probability Basics Review

## Setup:

- ▶ Random Experiment.  
Flip a fair coin twice.
- ▶ Probability Space.
  - ▶ **Sample Space:** Set of outcomes,  $\Omega$ .  
 $\Omega = \{HH, HT, TH, TT\}$   
(Note: **Not**  $\Omega = \{H, T\}$  with two picks!)
  - ▶ **Probability:**  $Pr[\omega]$  for all  $\omega \in \Omega$ .  
 $Pr[HH] = \dots = Pr[TT] = 1/4$ 
    1.  $0 \leq Pr[\omega] \leq 1$ .
    2.  $\sum_{\omega \in \Omega} Pr[\omega] = 1$ .
  - ▶ **Events.**  
Event  $A \subseteq \Omega$ ,  $Pr[A] = \sum_{\omega \in A} Pr[\omega]$ .

## Events:Poll

The following are events in the sample space corresponding to flipping a coin twenty times.

- (A) The first coin is a heads.
- (B) The last coin is a heads.
- (C) The outcome where every coin is a heads.
- (D) 7 out of 20 coins are heads.
- (E) The probability of all heads is  $1/2^{20}$ .

A, B, C, D

# Probability is Additive

## Theorem

(a) If events  $A$  and  $B$  are disjoint, i.e.,  $A \cap B = \emptyset$ , then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events  $A_1, \dots, A_n$  are pairwise disjoint, i.e.,  $A_k \cap A_m = \emptyset, \forall k \neq m$ , then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

## Proof:

(a)  $Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega]$   
 $= \sum_{\omega \in A} Pr[\omega] + \sum_{\omega \in B} Pr[\omega]$  since  $A \cap B = \emptyset$ .  $= Pr[A] + Pr[B]$

(b) Either induction, or argue over sample points.

# Consequences of Additivity

## Theorem

$$(a) \Pr[A \cup B] = \Pr[A] + \Pr[B] - \Pr[A \cap B];$$

(inclusion-exclusion property)

$$(b) \Pr[A_1 \cup \dots \cup A_n] \leq \Pr[A_1] + \dots + \Pr[A_n];$$

(union bound)

(c) If  $A_1, \dots, A_N$  are a **partition** of  $\Omega$ , i.e.,

pairwise disjoint and  $\bigcup_{m=1}^N A_m = \Omega$ , then

$$\Pr[B] = \Pr[B \cap A_1] + \dots + \Pr[B \cap A_N].$$

(law of total probability)


## Proof:

(b) follows from the fact that every  $\omega \in A_1 \cup \dots \cup A_n$  is included at least once in the right hand side.

Proofs for (a) and (c)? Next...

# Inclusion/Exclusion

$$Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$$



$$Pr[A] = x + y$$


$$Pr[B] = y + z$$

$$Pr[A \cap B] = y$$

$$Pr[A \cup B] = x + y + z$$

Another view. Any  $\omega \in A \cup B$  is in  $A \cap \bar{B}$ ,  $A \cup B$ , or  $\bar{A} \cap B$ . So, add it up.

## Roll a Red and a Blue Die.



$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$

$E_1$  = 'Red die shows 6';  $E_2$  = 'Blue die shows 6'


$E_1 \cup E_2$  = 'At least one die shows 6'

$$Pr[E_1] = \frac{6}{36}, Pr[E_2] = \frac{6}{36}, Pr[E_1 \cup E_2] = \frac{11}{36}.$$



# Total probability

Assume that  $\Omega$  is the union of the disjoint sets  $A_1, \dots, A_N$ .



Then,

$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$

Indeed,  $B$  is the union of the disjoint sets  $A_n \cap B$  for  $n = 1, \dots, N$ .

In “math”:  $\omega \in B$  is in exactly one of  $A_i \cap B$ .

Adding up probability of them, get  $Pr[\omega]$  in sum.


..Did I say...

Add it up.

# Conditional Probability.

**Definition:** The **conditional probability** of  $B$  given  $A$  is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



In  $A$ !

In  $B$ ?

Must be in  $A \cap B$ .

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$


## Conditional probability: example.

Two coin flips. First flip is heads. Probability of two heads?


$\Omega = \{HH, HT, TH, TT\}$ ; Uniform probability space.

Event  $A =$  first flip is heads:  $A = \{HH, HT\}$ .

$\Omega$  : uniform



New sample space:  $A$ ; uniform still.



Event  $B =$  two heads.

The probability of two heads if the first flip is heads.

**The probability of  $B$  given  $A$  is  $1/2$ .**


## A similar example.

Two coin flips. At least one of the flips is heads.


→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$ ; uniform.

Event  $A$  = at least one flip is heads.  $A = \{HH, HT, TH\}$ .



New sample space:  $A$ ; uniform still.




Event  $B$  = two heads.


The probability of two heads if at least one flip is heads.

**The probability of  $B$  given  $A$  is  $1/3$ .**

# Conditional Probability: A non-uniform example



Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red} | \text{Red or Green}] = \frac{3}{7} = \frac{Pr[\text{Red} \cap (\text{Red or Green})]}{Pr[\text{Red or Green}]}$$

## Another non-uniform example

Consider  $\Omega = \{1, 2, \dots, N\}$  with  $Pr[n] = p_n$ .

Let  $A = \{3, 4\}$ ,  $B = \{1, 2, 3\}$ .




$$Pr[A|B] = \frac{p_3}{p_1 + p_2 + p_3} = \frac{Pr[A \cap B]}{Pr[B]}.$$

## Yet another non-uniform example

Consider  $\Omega = \{1, 2, \dots, N\}$  with  $Pr[n] = p_n$ .

Let  $A = \{2, 3, 4\}$ ,  $B = \{1, 2, 3\}$ .




$$Pr[A|B] = \frac{p_2 + p_3}{p_1 + p_2 + p_3} = \frac{Pr[A \cap B]}{Pr[B]}.$$

# More fun with conditional probability.

Toss a red and a blue die, sum is 4,  
What is probability that red is 1?

$\Omega$  : Uniform



$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{3}; \text{ versus } Pr[B] = 1/6.$$


$B$  is more likely given  $A$ .



## Yet more fun with conditional probability.

Toss a red and a blue die, sum is 7,  
what is probability that red is 1?

$\Omega$  : Uniform



$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{6}; \text{ versus } Pr[B] = \frac{1}{6}.$$


Observing  $A$  does not change your mind about the likelihood of  $B$ .

## Such empty: poll

Suppose I toss 3 balls into 3 bins.

$A$  = "1st bin empty";  $B$  = "2nd bin empty."

$$\Omega = \{1, 2, 3\}^3$$



What is  $Pr[A|B]$ ?

- (A)  $1/27$
- (B)  $8/27$
- (C)  $1/8$
- (D) 0
- (E) 2


Next slide.

## Such empty..

Suppose I toss 3 balls into 3 bins.

$A$  = "1st bin empty";  $B$  = "2nd bin empty." What is  $Pr[A|B]$ ?

$$\Omega = \{1, 2, 3\}^3$$



$\omega = (\text{bin of red ball}, \text{bin of blue ball}, \text{bin of green ball})$

$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$

$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]} = \frac{(1/27)}{(8/27)} = 1/8; \text{ vs. } Pr[A] = \frac{8}{27}.$$

$A$  is less likely given  $B$ : If second bin is empty the first is more likely to have balls in it.

## Gambler's fallacy.

Flip a fair coin 51 times.

$A$  = “first 50 flips are heads”

$B$  = “the 51st is heads”

$Pr[B|A]$  ?

$A = \{HH \dots HT, HH \dots HH\}$

$B \cap A = \{HH \dots HH\}$

Uniform probability space.

$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{2}.$$

Same as  $Pr[B]$ .

The likelihood of 51st heads does not depend on the previous flips.

# Product Rule

Recall the definition:

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Hence,

$$Pr[A \cap B] = Pr[A]Pr[B|A].$$

Consequently,

$$\begin{aligned} Pr[A \cap B \cap C] &= Pr[(A \cap B) \cap C] \\ &= Pr[A \cap B]Pr[C|A \cap B] \\ &= Pr[A]Pr[B|A]Pr[C|A \cap B]. \end{aligned}$$

# Product Rule

## Theorem Product Rule

Let  $A_1, A_2, \dots, A_n$  be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

**Proof:** By induction.

Assume the result is true for  $n$ . (It holds for  $n = 2$ .) Then,

$$\begin{aligned} Pr[A_1 \cap \dots \cap A_n \cap A_{n+1}] &= Pr[A_1 \cap \dots \cap A_n]Pr[A_{n+1}|A_1 \cap \dots \cap A_n] \\ &= Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}]Pr[A_{n+1}|A_1 \cap \dots \cap A_n], \end{aligned}$$

so that the result holds for  $n + 1$ . □

# Correlation

An example.

Random experiment: Pick a person at random.

Event  $A$ : the person has lung cancer.

Event  $B$ : the person is a heavy smoker.

Fact:

$$Pr[A|B] = 1.17 \times Pr[A].$$

Conclusion:

- ▶ Smoking increases the probability of lung cancer by 17%.
- ▶ Smoking causes lung cancer.

# Correlation

Event  $A$ : the person has lung cancer. Event  $B$ : the person is a heavy smoker.  $Pr[A|B] = 1.17 \times Pr[A]$ .

A second look.

Note that

$$\begin{aligned} Pr[A|B] = 1.17 \times Pr[A] &\Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = 1.17 \times Pr[A] \\ &\Leftrightarrow Pr[A \cap B] = 1.17 \times Pr[A]Pr[B] \\ &\Leftrightarrow Pr[B|A] = 1.17 \times Pr[B]. \end{aligned}$$

Conclusion:

- ▶ Lung cancer increases the probability of smoking by 17%.
- ▶ Lung cancer causes smoking. Really?



# Causality vs. Correlation

Events  $A$  and  $B$  are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

$A$  and  $B$  being positively correlated does not mean that  $A$  causes  $B$  or that  $B$  causes  $A$ .

Other examples:

- ▶ Tesla owners are more likely to be rich. That does not mean that poor people should buy a Tesla to get rich.
- ▶ People who go to the opera are more likely to have a good career. That does not mean that going to the opera will improve your career.
- ▶ Rabbits eat more carrots and do not wear glasses. Are carrots good for eyesight?

# Proving Causality

Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).


Some difficulties:

- ▶  $A$  and  $B$  may be positively correlated because they have a common cause. (E.g., being a rabbit.)
- ▶ If  $B$  precedes  $A$ , then  $B$  is more likely to be the cause. (E.g., smoking.) However, they could have a common cause that induces  $B$  before  $A$ . (E.g., like math, CS70, Tesla.)

More about such questions later. For fun, check “N. Taleb: Fooled by randomness.”

# Total probability with Conditional Probability.

Assume that  $\Omega$  is the union of the disjoint sets  $A_1, \dots, A_N$ .



Then,


$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$

Indeed,  $B$  is the union of the disjoint sets  $A_n \cap B$  for  $n = 1, \dots, N$ . Thus,

$$Pr[B] = Pr[A_1]Pr[B|A_1] + \dots + Pr[A_N]Pr[B|A_N].$$

# Total probability

Assume that  $\Omega$  is the union of the disjoint sets  $A_1, \dots, A_N$ .



$$Pr[B] = Pr[A_1]Pr[B|A_1] + \dots + Pr[A_N]Pr[B|A_N].$$

## Simple Bayes Rule.

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}, Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

$$Pr[A \cap B] = Pr[A|B]Pr[B] = Pr[B|A]Pr[A].$$

$$\text{Bayes Rule: } Pr[A|B] = \frac{Pr[B|A]Pr[A]}{Pr[B]}.$$

## Is your coin loaded?

Your coin is fair w.p.  $1/2$  or such that  $Pr[H] = 0.6$ , otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

### Analysis:

$A =$  'coin is fair',  $B =$  'outcome is heads'

We want to calculate  $P[A|B]$ .

We know  $P[B|A] = 1/2$ ,  $P[B|\bar{A}] = 0.6$ ,  $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$\begin{aligned} Pr[B] &= Pr[A \cap B] + Pr[\bar{A} \cap B] = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}] \\ &= (1/2)(1/2) + (1/2)0.6 = 0.55. \end{aligned}$$

Thus,

$$Pr[A|B] = \frac{Pr[A]Pr[B|A]}{Pr[B]} = \frac{(1/2)(1/2)}{(1/2)(1/2) + (1/2)0.6} \approx 0.45.$$

## Independence:poll

**Definition:** Two events  $A$  and  $B$  are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Which Examples are independent?

- (A) Roll two dice,  $A$  = sum is 7 and  $B$  = red die is 1.
  - (B) Roll two dice,  $A$  = sum is 3 and  $B$  = red die is 1.
  - (C) Flip two coins,  $A$  = coin 1 is heads and  $B$  = coin 2 is tails.
  - (D) Throw 3 balls into 3 bins,  $A$  = bin 1 is empty and  $B$  = bin 2 is empty
- (A) and (C).

# Independence

**Definition:** Two events  $A$  and  $B$  are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice,  $A =$  sum is 7 and  $B =$  red die is 1 are independent;
- ▶ When rolling two dice,  $A =$  sum is 3 and  $B =$  red die is 1 are **not** independent;
- ▶ When flipping coins,  $A =$  coin 1 yields heads and  $B =$  coin 2 yields tails are independent;
- ▶ When throwing 3 balls into 3 bins,  $A =$  bin 1 is empty and  $B =$  bin 2 is empty are **not** independent;



## Independence: equivalent definition.

**Definition:** Two events  $A$  and  $B$  are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice,  $A = \text{sum is 7}$  and  $B = \text{red die is 1}$  are independent;  $Pr[A \cap B] = \frac{1}{36}$ ,  $Pr[A]Pr[B] = \left(\frac{1}{6}\right)\left(\frac{1}{6}\right)$ .
- ▶ When rolling two dice,  $A = \text{sum is 3}$  and  $B = \text{red die is 1}$  are **not** independent;  $Pr[A \cap B] = \frac{1}{36}$ ,  $Pr[A]Pr[B] = \left(\frac{2}{36}\right)\left(\frac{1}{6}\right)$ .
- ▶ When flipping coins,  $A = \text{coin 1 yields heads}$  and  $B = \text{coin 2 yields tails}$  are independent;  $Pr[A \cap B] = \frac{1}{4}$ ,  $Pr[A]Pr[B] = \left(\frac{1}{2}\right)\left(\frac{1}{2}\right)$ .
- ▶ When throwing 3 balls into 3 bins,  $A = \text{bin 1 is empty}$  and  $B = \text{bin 2 is empty}$  are **not** independent;  $Pr[A \cap B] = \frac{1}{27}$ ,  $Pr[A]Pr[B] = \left(\frac{8}{27}\right)\left(\frac{8}{27}\right)$ .

# Independence and conditional probability

**Fact:** Two events  $A$  and  $B$  are **independent** if and only if


$$Pr[A|B] = Pr[A].$$

Indeed:  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$ , so that

$$Pr[A|B] = Pr[A] \Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = Pr[A] \Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B].$$

# Conditional Probability: Pictures

Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$ . Note:  $Pr[B] \in (b_2, b_1)$ .
- ▶ Right:  $A$  and  $B$  are negatively correlated.  
 $Pr[B|A] = b_1 < Pr[B|\bar{A}] = b_2$ . Note:  $Pr[B] \in (b_1, b_2)$ .