

CS70 @ UC Berkeley, Spring 2026

Lecture 14 Counting I

March 5, 2026

Introduction

- The second half of the semester will focus on **probability theory**.
- In discrete probability, we will want to count the number of elements of a finite set.
- Counting such objects is important because, when all outcomes are equally likely, probabilities can be defined as

$$\Pr(A) = \frac{|A|}{|\Omega|}$$

where Ω is the set of possible **outcomes** and $A \subseteq \Omega$ is a subset of outcomes satisfying some property.


Introduction

- However, counting can quickly become **surprisingly subtle** and **difficult**.
- Many important objects in mathematics are **defined implicitly**.
- Seemingly small changes in constraints can dramatically change the count.
 - E.g., counting unconstrained **functions** $f: X \rightarrow S$ is relatively easy, but counting functions subject to certain restrictions is more involved.
 - Often we must count objects that satisfy several simultaneous constraints.
- **Direct counting is often infeasible**; e.g., some sets are extremely large.

Introduction

- Counting often requires **clever reasoning** rather than brute force.
- **Counting the same set in different ways** can reveal deep mathematical identities.
- **Techniques** include **bijections**, **inclusion-exclusion**, generating functions, and recursion. We will learn some of these techniques in this course.
- Developing good counting techniques is essential for studying discrete probability and many areas of computer science, statistics, and mathematics.

Sampling




inequivalent k -element objects depends on whether

- sampled objects are **ordered** or **unordered**
- sampling is done **with** or without **replacement**

Case 1: Ordered objects sampled with replacement

- #distinct polynomials over $\text{GF}(2)$ with degree ≤ 2 ?

$$P(x) = a_2x^2 + a_1x + a_0, \text{ where } a_0, a_1, a_2 \in \text{GF}(2)$$



$$S = \{0,1\}$$

Each path from the root to a leaf defines a unique polynomial over $\text{GF}(2)$


- #distinct polynomials over $\text{GF}(n)$ with degree $\leq k - 1$?

$$P(x) = a_{k-1}x^{k-1} + \dots + a_1x + a_0, \text{ where } a_0, a_1, \dots, a_{k-1} \in \text{GF}(n)$$

Answer: n^k

$$S = \{0,1, \dots, n - 1\}$$

Sampling




inequivalent k -element objects depends on whether

- sampled objects are **ordered** or **unordered**
- sampling is done with or **without replacement**

Case 2: Ordered objects sampled without replacement

ways to distribute 2 tasks to 3 compute nodes $\{A, B, C\}$, with **at most 1 task per node**. $S = \{A, B, C\}$



$3 \times 2 = 6$ leaves

General case: # ways to distribute k tasks to n compute nodes, with **at most 1 task per node**.

distinct length- k strings

$$= n(n-1)(n-2) \cdots (n-k+1)$$

$$= \frac{n!}{(n-k)!} \quad \text{“}k\text{th falling factorial of }n\text{”}$$

If $k = n$, then this expression becomes $n!$, which is the number of **permutations** on n distinct elements.

First Rule of Counting


Suppose an ordered object is formed by a succession of k choices where there are

n_1 possibilities for the 1st choice,
 n_2 possibilities for the 2nd choice,
⋮
 n_k possibilities for the k^{th} choice.

Then, the total number of distinct ordered objects is given by

$$n_1 \times n_2 \times \cdots \times n_k.$$

Sampling



inequivalent k -element objects depends on whether

- sampled objects are **ordered** or **unordered**
- sampling is done with or **without** **replacement**

Counting functions between two finite sets

$$f: X \rightarrow S, \text{ where } |X| = k \text{ and } |S| = n$$

The number of inequivalent functions f depends on:

- **Restrictions** on f
 - f is arbitrary (no restriction).
 - f is injective (one-to-one). Need $k \leq n$.
 - f is surjective (onto). Need $k \geq n$.
- Whether the elements of X and S are **distinguishable** or **indistinguishable**.

Case 1: f arbitrary; both X and S contain distinguishable elements

Case 2: f injective; both X and S contain distinguishable elements

Case 1 vs. Case 2

- Case 1: n^k
 - k -element ordered objects sampled with replacement
 - **arbitrary** $f: X \rightarrow S$; both X and S contain distinguishable elements
- Case 2: $n(n-1)\cdots(n-k+1)$
 - k -element ordered objects sampled without replacement
 - **injective** $f: X \rightarrow S$; both X and S contain distinguishable elements

For any fixed k :
$$\lim_{n \rightarrow \infty} \frac{n(n-1)\cdots(n-k+1)}{n^k} = 1$$


When the sample size k is small compared to the population size n , sampling with and without replacement are similar. Most of the functions are injective for $k \ll n$.

For $k = n$:
$$\lim_{n \rightarrow \infty} \frac{n!}{n^n} = \lim_{n \rightarrow \infty} \frac{\overset{\text{Stirling's approximation of } n!}{\sqrt{2\pi n} \left(\frac{n}{e}\right)^n}}{n^n} = \lim_{n \rightarrow \infty} \sqrt{2\pi n} \frac{1}{e^n} = 0$$

Almost no functions are injective for large $k = n$.


Balls and Bins

$f: X \rightarrow S$, where $|X| = k$ and $|S| = n$




Elements (balls) of X are **distinguishable**
Elements (bins) of S are distinguishable

These two configurations are **inequivalent**



The above two configurations become **equivalent** if the labels on the balls get removed.



Elements (balls) of X are **indistinguishable**
Elements (bins) of S are distinguishable

Case 3: Unordered objects sampled without replacement

Injective $f: X \rightarrow S$; the elements (balls) in X are **indistinguishable** while the elements (bins) in S are **distinguishable**

- For every configuration with **unlabeled** balls, there correspond $k!$ configurations with **labeled** balls.
- Hence, the number of **injections** $f: X \rightarrow S$ where X contains **indistinguishable** elements while S contains **distinguishable** elements is given by

$$\frac{n(n-1) \cdots (n-k+1)}{k!} = \frac{n!}{k!(n-k)!} = \binom{n}{k} = \binom{n}{n-k}$$

Second Rule of Counting

- Suppose an object is formed by a succession of k choices, but the order does not matter.
- If there exists an m -to-1 **surjective** map

$$g: \{\text{ordered objects}\} \rightarrow \{\text{unordered objects}\}$$

(i.e., for every $x \in \{\text{unordered objects}\}$, $|g^{-1}(x)| = m$), then

$$|\{\text{unordered objects}\}| = \frac{|\{\text{ordered objects}\}|}{m}$$

- Count as if order matters and then divide by m .

Second Rule of Counting


- More generally, if there exists an m -to-1 **surjective** map

$$g: A \rightarrow B$$

(i.e., for every $x \in B$, $|g^{-1}(x)| = m$), then


$$|B| = \frac{|A|}{m}$$

Example 1



- In how many different orders can an octopus put on its socks and shoes on its 8 limbs?
- For humans there are 6 ways:
 1. (Left **sock**, Left **shoe**, Right **sock**, Right **shoe**)
 2. (Left **sock**, Right **sock**, Left **shoe**, Right **shoe**)
 3. (Left **sock**, Right **sock**, Right **shoe**, Left **shoe**)
 4. (Right **sock**, Left **sock**, Left **shoe**, Right **shoe**)
 5. (Right **sock**, Left **sock**, Right **shoe**, Left **shoe**)
 6. (Right **sock**, Right **shoe**, Left **sock**, Left **shoe**)

Example 1




Socks: X_1, X_2, \dots, X_8

Shoes: Z_1, Z_2, \dots, Z_8

1. # Ways to order these 16 items = **16!** (by the **First Rule of Counting**). Let A denote the set of these length-16 sequences
2. But, not every $s \in A$ satisfies the constraint that **sock X_i should appear before shoe Z_i** . Let B denote the subset of **valid sequences** in A .
3. Each $s \in A$ can be mapped (g) to a valid sequence in B by **swapping Z_i with X_i if Z_i appears before X_i** .
4. $g: A \rightarrow B$ is a many-to-one surjection. For every $v \in B$, $|g^{-1}(v)| = 2^8$.
5. Hence, by the **Second Rule of Counting**, we obtain $\frac{16!}{2^8}$ for the answer.

Case 4: Unordered objects sampled with replacement


Arbitrary $f: X \rightarrow S$; the elements (balls) in X are **indistinguishable** while the elements (bins) in S are **distinguishable**.



4 **unlabeled** balls into
5 **labeled** bins,
no restriction on occupancy

Case 4: Unordered objects sampled with replacement

Arbitrary $f: X \rightarrow S$; the elements (balls) in X are **indistinguishable** while the elements (bins) in S are **distinguishable**.




k unlabeled balls into n labeled bins, no restrictions



Bijection

k unlabeled balls into $n + k - 1$ labeled bins, with ≤ 1 ball per bin

the length of consecutive occupied bins




Zeroth Rule of Counting

- If there exists a bijection (one-to-one correspondence) between two finite sets A and B , then $|A| = |B|$.

Case 4: Unordered objects sampled with replacement

Arbitrary $f: X \rightarrow S$; the elements (balls) in X are **indistinguishable** while the elements (bins) in S are **distinguishable**



Sometimes this representation is referred to “**stars and bars**”, where “stars” correspond to balls and “bars” to empty bins.

From **Case 3**, we know how to count the number of configurations for k **unlabeled balls** and $n + k - 1$ **labeled bins**, with **at most one ball per bin**:

$$\binom{n + k - 1}{k} = \binom{n + k - 1}{n - 1}$$

Example 2

Let n and k be positive integers. How many solutions does the following equation admit?

$$x_1 + x_2 + \cdots + x_n = k, \text{ where } x_i \in \mathbb{N}.$$

Answer: $\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$ k unlabeled balls into n labeled bins, no restrictions

Let n and $k > n$ be positive integers. How many solutions does the following equation admit?

$$x_1 + x_2 + \cdots + x_n = k, \text{ where } x_i \in \mathbb{N} \setminus \{0\}.$$

$$x_1 + x_2 + \cdots + x_n = k - n, \text{ where } x_i \in \mathbb{N}.$$

Answer: $\binom{k-1}{k-n} = \binom{k-1}{n-1}$ There exists a bijection between the solutions to these two equations.

Summary

$f: X \rightarrow S$, where $|X| = k$ and $|S| = n$

Sampling with replacement Sampling without replacement

	Elements of X	Elements of S	Arbitrary f	Injective f
Ordered	Distinguishable	Distinguishable	Case 1 n^k	Case 2 $\frac{n!}{(n-k)!}$
Unordered	Indistinguishable	Distinguishable	Case 4 $\binom{n+k-1}{k}$	Case 3 $\binom{n}{k}$