

# CS70 @ UC Berkeley, Spring 2026


## Lecture 25

### Continuous Probability Distribution II


April 23, 2026

# Random Breaks to Form a Triangle


- Suppose Alice samples two points independently and uniformly at random from the interval  $[0,1]$ , thereby obtaining 3 segments.



- What is the Probability that the three segments can form a triangle?



# Random Breaks to Form a Triangle




$$L_1 + L_2 + L_3 = 1$$

# Cumulative Distribution Function (CDF)


**Definition (CDF).** Given a random variable  $X$ , its cumulative distribution function  $F_X$  is defined as

$$F_X(a) = \mathbb{P}(X \leq a), \text{ for } a \in (-\infty, +\infty)$$

**Example:** Toss a fair coin twice.  
 $X(\omega) = \text{Heads in } \omega \in \Omega.$



$U \sim \text{Uniform}[0, L]$   
Continuous CDF



# Convergence in Distribution

CDF plays a central role in probability theory.

**Definition (Convergence in distribution):**  $X_n \xrightarrow{d} X$  as  $n \rightarrow \infty$

A sequence of random variables  $X_1, X_2, X_3, \dots$  is said to converge to another random variable  $X$  in distribution if

$$\lim_{n \rightarrow \infty} F_{X_n}(x) = F_X(x),$$

for all  $x \in \mathbb{R}$  where  $F_X(x)$  is continuous.

## Examples:

1.  $X_n \sim \text{Binomial}(n, p_n)$ , where  $p_n = \lambda/n$ . Then,
2.  $Y_n \sim \text{Geometric}(p_n)$ , where  $p_n = \lambda/n$ . Then,


# Normal Distribution

Definition (Normal Random Variable):  $X \sim \text{Normal}(\mu, \sigma^2)$

A continuous random variable  $X$  with probability density function

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/(2\sigma^2)},$$

where  $\mu \in \mathbb{R}$  and  $\sigma^2 > 0$ .



The density becomes more spread out as  $\sigma^2$  increases, while  $\mu$  determines where the center is.

# Normal Distribution

Definition (Normal Random Variable):  $X \sim \text{Normal}(\mu, \sigma^2)$

A continuous random variable  $X$  with probability density function

$$f_X(x) = \frac{1}{2\pi\sigma^2} e^{-(x-\mu)^2/(2\sigma^2)},$$


where  $\mu \in \mathbb{R}$  and  $\sigma^2 > 0$ .

1.  $\int_{-\infty}^{\infty} f_X(x) dx = 1$
2.  $\mathbb{E}[X] = \mu$
3.  $\text{Var}[X] = \sigma^2$

**Proof of 2:**

# CDF of the Standard Normal Distribution


$\Phi(x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$  denotes the c.d.f. of Normal(0,1).



# Sum of Independent Normal Random Variables


Suppose  $X, Y \sim \text{Normal}(0,1)$  are **independent**.

$$f_{X,Y}(x, y) = f_X(x)f_Y(y) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \frac{1}{\sqrt{2\pi}} e^{-y^2/2} = \frac{1}{2\pi} e^{-(x^2+y^2)/2}$$



$$\begin{aligned} X_\theta &= X \cos \theta + Y \sin \theta \\ Y_\theta &= -X \sin \theta + Y \cos \theta \end{aligned}$$

How are these distributed?



# Sum of Independent Normal Random Variables

Suppose  $X, Y \sim \text{Normal}(0,1)$  are **independent**.

$$f_{X,Y}(x, y) = f_X(x)f_Y(y) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \frac{1}{\sqrt{2\pi}} e^{-y^2/2} = \frac{1}{2\pi} e^{-(x^2+y^2)/2}$$

$$\begin{aligned} X_\theta &= X \cos \theta + Y \sin \theta \\ Y_\theta &= -X \sin \theta + Y \cos \theta \end{aligned}$$

Rotationally symmetric  $\Rightarrow$

$X_\theta, Y_\theta \sim \text{Normal}(0,1)$  and they are **independent**

# Fun Fact about the Normal Distribution

Let  $X$  and  $Y$  be arbitrary random variables on the same probability space.

$$\begin{pmatrix} X_\theta \\ Y_\theta \end{pmatrix} = \underbrace{\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}}_M \begin{pmatrix} X \\ Y \end{pmatrix} \quad \begin{array}{l} M^T M = I \\ M M^T = I \end{array} \quad M \text{ is an orthogonal matrix}$$

$$X_\theta^2 + Y_\theta^2 = (X_\theta \ Y_\theta) \begin{pmatrix} X_\theta \\ Y_\theta \end{pmatrix} = (X \ Y) M^T M \begin{pmatrix} X \\ Y \end{pmatrix} = (X \ Y) \begin{pmatrix} X \\ Y \end{pmatrix} = X^2 + Y^2$$

⇒ Every orthogonal transformation of a vector preserves its length.

# Central Limit Theorem (CLT)

**Theorem (Weak Law of Large Numbers, Lecture 23).** Let  $X_1, X_2, X_3, \dots$  be a sequence of i.i.d. random variables with **finite mean**  $\mu$  and **finite variance**  $\sigma^2$ . Let  $S_n = X_1 + \dots + X_n$ . Then,  $\lim_{n \rightarrow \infty} \mathbb{P} \left( \left| \frac{S_n}{n} - \mu \right| > \varepsilon \right) = 0, \forall \varepsilon > 0$ .

WLLN is a statement about the convergence of the sample average  $\frac{S_n}{n}$  to its mean as  $n \rightarrow \infty$

**Theorem (CLT):** Let  $X_1, X_2, X_3, \dots$  be a sequence of i.i.d. random variables with **finite mean**  $\mu$  and **finite variance**  $\sigma^2$ . Let  $S_n = X_1 + \dots + X_n$ . Then,

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[ \frac{\sqrt{n}}{\sigma} \left( \frac{S_n}{n} - \mu \right) \leq x \right] = \Phi(x), \quad \forall x \in \mathbb{R},$$


where  $\Phi(x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$  denote the c.d.f. of Normal(0,1).

$\mathbb{E} \left[ \frac{S_n}{n} \right] = \mu$  and  $\text{Var} \left( \frac{S_n}{n} \right) = \frac{\sigma^2}{n}$ . Formally, we write  $\frac{\sqrt{n}}{\sigma} \left( \frac{S_n}{n} - \mu \right) \xrightarrow{d} X \sim \text{Normal}(0,1)$  as  $n \rightarrow \infty$ .


CTL is a statement about the **scaled fluctuation** of  $\frac{S_n}{n}$  about  $\mu$  as  $n \rightarrow \infty$ .

**Application:** For large  $n$ ,  $\frac{S_n}{n}$  is well approximated by Normal( $\mu, \frac{\sigma^2}{n}$ )

# CLT Demo



# CLT Demo







# Can you now explain this behavior?

$N \sim \text{Poisson}(\lambda)$ , where intensity  $\lambda > 0$ .

$$\mathbb{P}(N = k) = e^{-\lambda} \frac{\lambda^k}{k!}, \text{ for } k \in \mathbb{N}.$$

- # rain drops hitting a surface per second
- # radioactive particles emitted by radioactive material during an interval of time




Looks more and more like a “bell curve” as  $\lambda$  gets large. We will see why this happens.

# Poisson and CTL


$Y \sim \text{Poisson}(5000)$

$n = 1$




$X_1, \dots, X_{5000} \stackrel{\text{i.i.d.}}{\sim} \text{Poisson}(1)$

$S_n = X_1 + \dots + X_n \quad n = 5000$



- Recall that if  $Z_1 \sim \text{Poisson}(\lambda_1), \dots, Z_n \sim \text{Poisson}(\lambda_n)$  are independent, then  $Z_1 + \dots + Z_n \sim \text{Poisson}(\lambda_1 + \dots + \lambda_n)$
- So, in the above example,  $Y$  and  $X_1 + \dots + X_n$  are identically distributed.
- However, CLT implies  $\sqrt{n} \left( \frac{S_n}{n} - 1 \right)$  converges to  $\text{Normal}(0,1)$  as  $n \rightarrow \infty$ .


# How to estimate $\pi$ using a needle




The 18<sup>th</sup> century method  
(due to Buffon)

Assume an infinite array of parallel lines with gap size  $d$ . If a needle of length  $\ell \leq d$  is thrown uniformly at random, what is the probability that the needle intersects the grid?

Hit




Miss




# How to estimate $\pi$ using a needle

$$Y \sim \text{Uniform} \left[ 0, \frac{d}{2} \right] \quad \Theta \sim \text{Uniform} \left[ -\frac{\pi}{2}, \frac{\pi}{2} \right]$$



- Let  $Y$  be the distance between the center of the needle to the closest line.
- For a given  $\theta$ , the needle will intersect the grid **if and only if**  $y \leq \frac{\ell}{2} \cos \theta$ .



Let  $I_k$  be the indicator RV for the  $k$ th trial being successful, and let  $S_n = I_1 + \dots + I_n$ .


# How to estimate $\pi$ using a needle

$\hat{\pi}(I_1, \dots, I_n)$  from computer experiments for  $d = \ell$


Experiment	Sample size $n$		
	100	1,000	10,000
1	2.778	3.257	3.120
2	3.571	3.091	3.136
3	3.226	3.160	3.137
4	4.167	3.200	3.152


Variability decreases as  $n$  increases.

CLT can be used to approximate the uncertainty of the estimator for a given sample size  $n$ .




# An Alternate Solution to Buffon's Needle Problem


1. Suppose the needle consists of two pieces of lengths  $\ell_1$  and  $\ell_2$  
2.  $p_\ell := \mathbb{P}(\text{Needle of length } \ell \text{ intersects grid}) = p_{\ell_1} + p_{\ell_2}$ , which is satisfied if  $p_\ell = a\ell$ , for some constant  $a > 0$ .
3. Consider a polygonal chain consisting of  $n$  line segments of lengths  $\ell_1, \dots, \ell_n$ , where  $\sum_{k=1}^n \ell_k = \ell$  and  $\ell_k < d$  for all  $k = 1, \dots, n$ .



4. Let  $I_k$  be an indicator for the segment  $k$  intersecting the grid.
5. Let  $X$  = the total number of intersections. Then  $X = \sum_{k=1}^n I_k$  and  $\mathbb{E}[X] =$
6. A circle (infinitely many line segments) with diameter  $d$  (and hence length  $\ell = \pi d$ ) always intersects the grid **exactly twice**. So,  $\mathbb{E}[X] =$ .



# Convex Polygon Throws



Consider a **sufficiently small** (so it cannot cross more than one line at once) convex polygon with circumference  $C$ .

$X$  = the total number of intersections in a throw.